

Sporobolus africanus

System: Terrestrial

| Kingdom | Phylum | Class | Order | Family |
|---------|---------------|------------|-----------|---------|
| Plantae | Magnoliophyta | Liliopsida | Cyperales | Poaceae |

Common name Parramatta grass (English), rat's-tail grass (English), rat-tail grass (English), smut grass (English), African dropseed (English), tufty grass (English), motie hikutaha (Niuean), cat's tail (English), cape grass (English)

Synonym *Sporobolus indicus* var. *capensis*

Similar species

Summary The rat-tail grass *Sporobolus indicus* is a tuft forming perennial grass native to various countries in continental Africa and often introduced as a pasture grass. It is capable of forming dense stands once established which can exclude native species and alter ecological conditions such as light, nutrients and moisture availability. As a pasture weed, it is capable of having serious economic impacts and reducing land value. Management of *S. indicus* at present is difficult and costly due to its ease of spread and the creation of a very large seed bank that remains viable for up to 10 years.



[view this species on IUCN Red List](#)

Species Description

Sporobolus indicus is a perennial grass that grows in dense tufts (caespitose). Sheathes are papery and culms are erect being 30 - 110 cm long and 1.5 - 3 mm in diameter. Ligule a fringe of hairs. Leaf-blades flat, or convolute; 20– 40 cm long and 1 – 4 mm wide. Leaf-blade apex filiform. Inflorescence a contracted or spiciform, linear panicle 10 - 35 cm long, contracted about primary branches. Primary panicle branches appressed; 1 - 2 cm long; bearing spikelets almost to the base. The panicle branches glabrous in axils. Spikelets are solitary with fertile spikelets pedicelled. Spikelets comprising one fertile florets without rhachilla extension. Spikelets are lanceolate and subterete being (1.5-)2.1 - 2.8 mm long; breaking up at maturity; disarticulating below each fertile floret. Glumes are deciduous and dissimilar being shorter than the spikelet. The lower glume is oblong being 0.4 – 0.7 mm long about 0.4 - 0.5 of the length of the upper glume which is usually 1 – 1.5 mm long. Upper and lower glumes are without keels and lateral veins while the lower glume has an obtuse apex and the upper glume has an acute apex. Fertile lemma are ovate; (1.5-)2.1 – 2.8 mm long, membranous and dark green without a keel and 1-veined. The lemma apex is acute and the palea is 2-veined with approximate keels. Flowers have 3 anthers, 0.6 - 0.8 mm long. *Sporobolus indicus* produces an ellipsoid quadrangular caryopsis with a free, soft pericarp. Caryopsis is (0.8-) 1.1 - 1.2 mm long (Clayton *et al.*, 2006 onwards). Please follow this link [PIER, 2009](#) for images.

Notes

Much of the literature refers to *Sporobolus indicus* as its synonym, *S. africanus*.

Uses

In Mexico, the fruits are regarded as diuretic and antispasmodic. The fruit decoction is used to bathe wounds and heal sores in the mouth. A sirup prepared from the fruit is taken to overcome chronic diarrhea. The astringent bark decoction is a remedy for mange, ulcers, dysentery and for bloating caused by intestinal gas in infants. In the Philippines, the sap of the bark is used to treat stomatitis in infants. The juice of the fresh leaves is a remedy for thrush. A decoction of the leaves and bark is employed as a febrifuge. In southwestern Nigeria, an infusion of shredded leaves is valued for washing cuts, sores and burns. Researchers at the University of Ife have found that an aqueous extract of the leaves has antibacterial action, and an alcoholic extract is even more effective. The gum-resin of the tree is blended with pineapple or soursop juice for treating jaundice. Most of the other uses indicate that the fruits, leaves and bark are fairly rich in tannin (Global Compendium of Weeds, 2007).

Habitat Description

Sporobolus indicus is capable of naturalising in disturbed areas such as along roadsides and in grasslands and pastures in Hawaii (PIER, 2009). Other reports from New Zealand and Niue confirm that this is common in other areas where *S. indicus* has established (PIER, 2009). In Queensland, Australia *S. indicus* is a serious pasture weed which has significant economic impacts (Department of Primary Industries and Fisheries, 2007). Growth in pasture is maximised in the absence of ground cover as seedlings are susceptible to competitive effects when young (Department of Primary Industries and Fisheries, 2007).

Reproduction

Sporobolus indicus produces a large amount of seeds, up to 80 000 / m², with 90% of them being viable. Seed is produced all year in Australia, but mostly in summer and autumn. Seeding is difficult to prevent and quickly builds up a large seed bank which remains viable for up to 10 years. Seed germination occurs when there is low levels of competition from cover-forming species like pasture grasses, allowing *S. indicus* to take advantage of disturbance events. Early seedling survival is sensitive to competition until the seedlings are about 5 cm tall. At this stage, they are much more resistant to competition and moisture stress. In a pasture, *S. indicus* will not be selectively grazed, while the grazing of the more palatable pasture species allows more *S. indicus* seeds to germinate successfully. The adult plant is capable of living over four years, flowers within 3 months, and is capable of producing seed at any time given favourable conditions (Department of Primary Industries and Fisheries, 2007).

General Impacts

Sporobolus indicus is capable of dominating particular sites, forming dense cover when it becomes established which can result in the exclusion and decline of native species. This also has the effect of altering natural ecological conditions, such as light, nutrients and moisture availability (Gray *et al.*, 2005). *S. indicus* quite often forms exclusive ground covers with communities of other introduced species which has thought to have contributed to the decline of a number of threatened species (Cronk, 1986b; Gray *et al.*, 2005; 2009). As *S. indicus* provides a good combustible material, dense stands may be able to propagate wildfires which could enormously retard the recovery of native shrubs and herbs. (Loope *et al.*, 1992). When established in pastures, *S. indicus* can dramatically decrease producers' economic viability and lower land values, with current infestations in Australia costing the pastoral industry in the vicinity of AU\$ 60 million per year in lost production and control costs (Department of Primary Industries and Fisheries, 2007).



GLOBAL INVASIVE SPECIES DATABASE

FULL ACCOUNT FOR: *Sporobolus africanus*

Management Info

Please follow this link for detailed information on [the management and control of *Sporobolus indicus* \(rat-tail grass\)](#). A summary is provided below.

Preventative measures: Guidelines have been produced by the Department of Primary Industries and Fisheries (2007) outlining ways to prevent spread of rat-tail grass. These include cleaning vehicles and machinery as well as providing quarantine areas for livestock grazing in infested areas.

Physical control: While, hand pulling or digging is effective on individual plants, on a larger scale, this would be too labour intensive. Additionally, slashing promotes seed production and increases the risk of spreading seeds. If used correctly, fire can play a useful role in reducing the soil seed bank and encourage remaining seed to germinate further depleting seed reserves (Department of Primary Industries and Fisheries, 2007).

Chemical control: Both flupropanate and glyphosate herbicides have been identified as being effective for control of *S. indicus* in pastures, with application methods differing depending on the scale of infestation (Department of Primary Industries and Fisheries, 2007).

Cultural control: The Department of Primary Industries and Fisheries (2007) has suggested that the spread of *S. indicus* can be prevented by changing the way landowners manage their land, such as reducing bare or waste areas and preventing pasture overgrazing.

Biological control: Biological control of *S. indicus* was investigated using the stem wasp, *Tetramesa sp.* and the smut, *Ustilago sporoboli-indici*. Both were unsuccessful and future efforts are concentrated on the development of a pathogen already present in Australia as a mycoherbicide (Palmer *et al.*, 2010).

Integrated management: At present a combination of preventative measures, chemical and cultural control techniques is used to effectively control current infestations and prevent spread of *S. indicus*. The large persistent seed bank created by infestations however make it a labour intensive and expensive procedure (Department of Primary Industries and Fisheries, 2007).

Pathway

The seeds of *Sporobolus indicus* become sticky when wet, allowing them to stick to animals, clothes, vehicles and machinery (Department of Primary Industries and Fisheries, 2007). The seeds of *S. indicus* are also capable of spreading via soil on machinery and vehicles (Department of Employment, Economic Development and Innovation, 2010)The seeds of *Sporobolus indicus* are capable of spreading via soil on machinery and vehicles (Department of Employment, Economic Development and Innovation, 2010)The seeds of *Sporobolus indicus* can be propagated through contaminated hay or pasture seed (Department of Primary Industries and Fisheries, 2007).

Principal source: [Department of Primary Industries and Fisheries, 2007. Weedy *Sporobolus* grasses. Best Practice Manual](#). The original edition (1999, 2001) was compiled by the Giant Rats Tail team: Felicity McIntosh, Bill Schulke, Graeme Elphinstone, Steven Bray, Wayne Vogler, Col Paton, Bernie Shore, Michael Yee and Nigel Gallas. The revised edition (2007) has been updated by Steven Bray and David Officer.



GLOBAL INVASIVE SPECIES DATABASE

FULL ACCOUNT FOR: *Sporobolus africanus*

Compiler: IUCN SSC Invasive Species Specialist Group (ISSG) with support from the Overseas Territories Environmental Programme (OTEP) project XOT603, a joint project with the Cayman Islands Government - Department of Environment

Review: Under expert review

Publication date: 2010-06-02

ALIEN RANGE

[2] AUSTRALIA

[1] LESSER ANTILLES

[3] NEW ZEALAND

[1] NORFOLK ISLAND

[1] PITCAIRN

[9] UNITED STATES

[5] FRENCH POLYNESIA

[1] MAURITIUS

[1] NIUE

[1] NORTHERN MARIANA ISLANDS

[3] SAINT HELENA

BIBLIOGRAPHY

26 references found for *Sporobolus africanus*

Management information

Bourd?t, G. W. , Fowler, S. V. , Edwards, G. R. , Kriticos, D. J. , Kean, J. M. , Rahman, A. and Parsons, A. J., 2007. Pastoral weeds in New Zealand: Status and potential solutions , New Zealand Journal of Agricultural Research, 50: 2, 139 ? 161

[Department of Employment, Economic Development and Innovation, Queensland, 2009. Biological control of weedy *Sporobolus grasses*](#)

Summary: Available from: http://www.dpi.qld.gov.au/4790_13778.htm [Accessed 10 April 2010]

[Department of Employment, Economic Development and Innovation, Queensland, 2010. Parramatta grass](#)

Summary: Available from: http://www.dpi.qld.gov.au/4790_7333.htm [Accessed 10 April 2010]

[Department of Primary Industries and Fisheries, 2007. Weedy *Sporobolus grasses*. Best Practice Manual. The original edition \(1999, 2001\) was compiled by the Giant Rats Tail team: Felicity McIntosh, Bill Schulke, Graeme Elphinstone, Steven Bray, Wayne Vogler, Col Paton, Bernie Shore, Michael Yee and Nigel Gallas. The revised edition \(2007\) has been updated by Steven Bray and David Officer.](#)

Summary: Available from:

http://www.dpi.qld.gov.au/documents/Biosecurity_GeneralPlantHealthPestsDiseaseAndWeeds/Weedy-Sporobolus-manual.pdf [Accessed 10 April 2010]

Gray, Alan; Paul David Robinson and Stedson Stroud, 2009. Use of the Biological Flora framework in the United Kingdom Overseas Territories: *Euphorbia organoides* L. Biological Conservation 142 (2009) 1754?1766

[IUCN/SSC Invasive Species Specialist Group \(ISSG\).. 2010. A Compilation of Information Sources for Conservation Managers.](#)

Summary: This compilation of information sources can be sorted on keywords for example: Baits & Lures, Non Target Species, Eradication, Monitoring, Risk Assessment, Weeds, Herbicides etc. This compilation is at present in Excel format, this will be web-enabled as a searchable database shortly. This version of the database has been developed by the IUCN SSC ISSG as part of an Overseas Territories Environmental Programme funded project XOT603 in partnership with the Cayman Islands Government - Department of Environment. The compilation is a work under progress, the ISSG will manage, maintain and enhance the database with current and newly published information, reports, journal articles etc.

[Loope, Lloyd L., Ronald J. Nagata, and Arthur C. Medeiros, 1992. Alien Plants in Haleakala National Park.](#)

Summary: Available from: <http://www.hear.org/books/apineh1992/pdfs/apineh1992V5loopeetal.pdf> [Accessed 10 April 2010]

[Pacific Islands Ecosystems at Risk \(PIER\), 2009. *Sporobolus africanus* \(Poir.\) Robyns & Tournay, Poaceae](#)

Summary: Available from: http://www.hear.org/pier/species/sporobolus_africanus.htm [Accessed 10 April 2010]



GLOBAL INVASIVE SPECIES DATABASE

FULL ACCOUNT FOR: *Sporobolus africanus*

[Palmer, W.A. 2008. Biological control of weedy *Sporobolus* grasses by two host specific agents. Final report. Meat & Livestock Australia Limited, North Sydney. 63 pp.](#)

Summary: Available from:

<http://www.mla.com.au/TopicHierarchy/ResearchAndDevelopment/ResearchAndDevelopmentDetails.htm?projectId=1520> [Accessed 10 April 2010]

Palmer, W.A., T.A. Heard, & A.W. Sheppard, 2010. A review of Australian classical biological control of weeds programs and research activities over the past 12 years. *Biological Control* 52 (2010) 271-287

Palmer, W.A., Yobo, K.S. and Witt, A.B.R. 2008. Prospects for the biological control of the weedy *Sporobolus* grasses in Australia. In: Proceedings of the 16th Australian Weeds Conference. R.D. Van Klinken, V.A. Osten, F.D. Panetta and J.C. Scanlan, eds. Queensland Weeds Society, Brisbane. pp. 264-266.

Shrestha Sangita; Adkins Stephen W.; Graham Glenn C.; Loch Donald S., 2005. An identification tool for the Australian weedy *Sporobolus* species based on random amplified polymorphic DNA (RAPD) profiles. *Australian journal of agricultural research* 2005, vol. 56, no2, pp. 157-167

[UK Overseas Territories Conservation Forum \(UKOTCF\). 2010. Managing alien plants on the outer islands of Tristan da Cunha \(OTEP TDC403\)](#)

Summary: Available from: <http://www.ukotcf.org/infoDB/infoSourcesDetail2.cfm?module=projects&refID=213> [Accessed 10 April 2010]

Witt A.B.R. and A.J. McConnachie, 2003. The potential for classical biological control of invasive grass species with special reference to invasive *Sporobolus* spp. (Poaceae) in Australia IN Proceedings of the XI International Symposium on Biological Control of Weeds Canberra, Australia, 27 April-2 May 2003 Edited by J.M. Cullen, D.T. Briese, D.J. Kriticos, W.M. Lonsdale, L. Morin and J.K. Scott

Yobo, K.S., Laing, M.D., Palmer, W.A. and Shivas, R.G. 2009. Evaluation of *Ustilago sporoboli-indici* as a classical biological control agent for invasive *Sporobolus* grasses in Australia. *Biological Control* 50(1): 7-12.

General information

Aplet, G. H., S. J. Anderson & C. P. Stone, 1991. Association between feral pig disturbance and the composition of some alien plant assemblages in Hawaii Volcanoes National Park. *Vegetatio* 95: 55-62, 1991.

[Clayton, W.D., Harman, K.T. and Williamson, H. 2006 onwards. GrassBase - The Online World Grass Flora. *Sporobolus africanus*](#)

Summary: Available from: <http://www.kew.org/data/grasses-db/www/imp09589.htm> Accessed 10 April 2010]

Cronk, Q. C. B., 1986a. The Decline of the St Helena Ebony *Trochetiopsis melanoxyton*. *Biological Conservation* 35 (1986) 159-172

Cronk, Q. C. B., 1986b. The decline of the St Helena gumwood *Commidendrum robustum*. *Biological Conservation* Volume 35, Issue 2, 1986, Pages 173-186

Cronk, Q. C. B., 1989. The Past and Present Vegetation of St Helena. *Journal of Biogeography*, Vol. 16, No. 1 (Jan., 1989), pp. 47-64 (article consists of 18 pages)

[Global Compendium of Weeds \(GCW\), 2007. *Sporobolus africanus* \(Poaceae\)](#)

Summary: Available from: http://www.hear.org/gcw/species/sporobolus_africanus/ [Accessed 10 April 2010]

Gray, Alan; Tara Pelembe and Stedson Stroud, 2005. The conservation of the endemic vascular flora of Ascension Island and threats from alien species. *Oryx* (2005), 39:4:449-453 Cambridge University Press

[Integrated Taxonomic Information System \(ITIS\), 2010. *Sporobolus indicus* var. *capensis* Engelm.](#)

Summary: Available from: http://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=566313 [Accessed 10 April 2010]

[Oldfield, S., Procter, D., & Fleming L.V. \(1999\). Biodiversity: the UK Overseas Territories. Joint Nature Conservation Committee.](#)

Summary: Available from:

http://www.widecast.org/Resources/Docs/Procter_and_Fleming_1999_Biodiversity_in_UKOTs.pdf [Accessed 13 April 2010]

[USDA-ARS, 2010. Taxon: *Sporobolus indicus* \(L.\) R. Br. var. *capensis* Engl. National Genetic Resources Program. Germplasm Resources Information Network - \(GRIN\) \[Online Database\]. National Germplasm Resources Laboratory, Beltsville, Maryland.](#)

Summary: Available from: http://www.ars-grin.gov/cgi-bin/npgs/html/tax_search.pl?Sporobolus%20africanus [Accessed 10 April 2010]

[USDA-NRCS, 2010. *Sporobolus indicus* \(L.\) R. Br. var. *capensis* Engelm. rat-tail grass The PLANTS Database \(<http://plants.usda.gov>, 10 April 2010\). National Plant Data Center, Baton Rouge, LA 70874-4490 USA.](#)

Summary: Available from: <http://plants.usda.gov/java/profile?symbol=SPINC3> [Accessed 10 April 2010]