

MN (Minor) *Alectoris chukar*

Date assessed	2021-09-21
Year published	2022
Eicat category	MN (Minor)
Justification for EICAT assessment	The Chukar feeds on native invertebrates species in the U.S.A.. No population level impacts have been observed (Cole et al. 1995; Larsen et al. 2007).The Chukar hybridises with the Rock partridge and Red-legged partridge in Europe in the wild. No population level impacts have been observed. (Barilani et al. 2007a; Barilani et al. 2007b; Casas et al. 2012; Blanco-Aguilar et al. 2008; Barbanera et al. 2011; Barbanera et al. 2009; Barbanera 2005; Tejedor et al. 2007).
Confidence rating	Low
Mechanism(s) of maximum impact	Predation; Hybridisation
Countries of most severe impact	U.S.A.; France; Greece; Italy; Spain; Portugal
Description of impact	The Chukar hybridises with the Rock partridge and Red-legged partridge in Europe in the wild. It also feeds on native invertebrate and plant species.
Assessor	Thomas Evans; Tom Allmert
Contributors	
Reviewers	EICAT authority
Recommended citation	Thomas Evans; Tom Allmert. (2024). <i>Alectoris chukar</i> . IUCN Environmental Impact Classification for Alien Taxa (EICAT) .

