

**Caiman crocodilus**

**System:** Freshwater\_terrestrial

Kingdom	Phylum	Class	Order	Family
Animalia	Chordata	Reptilia	Crocodilia	Alligatoridae

<b>Common name</b>	polulo (English, Mexico), tinga (Spanish), cocodrilo (English, Mexico), lagarto (Spanish), ocoroche (English, Mexico), punamnah (Palikur, Brazil), yacaré (English, Peru), baba (English, Venezuela), guajipal (English, Nicaragua), common caiman (English, Puerto Rico), tulisio (English, Mexico), spectacled caiman (English), jacaretinga (Portuguese, Brazil), cascarudo (Spanish), babilla (English, Colombia), caiman de Brasil (English, Brazil), babiche (Spanish), caimán de anteojos (Spanish, Venezuela, Spain), caimán sudamericano (Spanish), cachirré (English, Mexico), jacaré-tinga (Portuguese, Brazil), caiman blanco (English, French Guiana), lagarto blanco (English, Peru), caimán (Spanish), yacaré blanco (Spanish)
<b>Synonym</b>	<i>Lacerta crocodilus</i> , (Linnaeus, 1758) <i>Caiman sclerops</i> , (Schneider, 1801) <i>Crocodilus sclerops</i> , Schneider 1801 (fide Wwemuth & Mertens 1977) <i>Alligator sclerops</i> , Dumeril & Bibron 1836: 79 <i>Caiman crocodilus</i> , Conant & Collins 1991: 40 <i>Caiman crocodilus</i> , Schwartz & Henderson 1991: 666 <i>Caiman crocodilus</i> , Gorzula & Senaris 1999 <i>Caiman crocodylus</i> , Lehr 2002: 69 <i>Caiman crocodilus apaporiensis</i> , Medem 1955 <i>Caiman sclerops apaporiensis</i> , Medem 1955 <i>Caiman crocodilus apaporiensis</i> , Nickel & Auliya 2004 <i>Caiman crocodilus chiapasius</i> , (Bocourt 1876) <i>Alligator (Jacare) chiapasius</i> , Bocourt 1876 <i>Caiman crocodilus fuscus</i> , (Cope 1868) <i>Perosuchus fuscus</i> , Cope 1868 <i>Perosuchus fuscus</i> , Gray 1869: 171 <i>Caiman crocodilus fuscus</i> , Nickel & Auliya 2004
<b>Similar species</b>	<i>Caiman latirostris</i> , <i>Caiman yacare</i>

<b>Summary</b>	The common caiman ( <i>Caiman crocodilus</i> ), is currently the most abundant crocodilian species and is the most harvested crocodile in the hide industry. Native to South and Central America, <i>C. crocodilus</i> has been introduced and has established in America, Puerto Rico, and Cuba. It poses a threat to native crocodilians through competition and is believed to be responsible for the introduction of the exotic parasite known as "caiman tongueworm" which infects local fish species in Puerto Rico.
----------------	--



[view this species on IUCN Red List](#)

## Species Description

The common caiman is a relatively small to medium sized crocodilian. Males generally reach 2.0-2.5 m with the largest specimens reported to approaching 3 m. Females are smaller, reaching a mean maximum size of 1.4 m, with larger specimens reaching 1.8 m. The common caiman is also known as the spectacled caiman, due to a bony ridge present between the eyes (infra-orbital bridge) which appears to join the eyes like a pair of spectacles. Juveniles are brown-cream colored with black spots and bands on the body and tail. Adults are dull olive-green to black with black bands basically on the tail. The enlarged fourth tooth of the lower jaw is not visible when the jaws are closed, as it is in all true crocodiles species (Behler, 1979), but in the oldest individuals, the front teeth and the fourth tooth of the lower jaw can perforate the upper jaw and can be visible when the mouth is closed.

## Notes

Four or five subspecies of *C. crocodilus* are recognized: *C. c. crocodilus* which is prevalent throughout Venezuela and the Amazon from Colombia to Brazil; *C. c. chiapasius* which inhabits most of Central America from south Mexico to the pacific coast of Colombia; *C. c. fuscus* which inhabits Atlantic coastal drainages of Colombia to western Venezuela; and *C. c. apaporiensis*, a narrower snouted form which is thought to only inhabit the upper Apaporis River in Colombia and is believed in need of conservation effort (CSG, 2008; Ross 1998). Some authorities believe that *C. yacare* is not a full species, but a subspecies (*C. c. yacare*) that occurs in southern Brazil, Bolivia, Paraguay and northern Argentina. All subspecies of *C. crocodilus* are in CITES Appendix II, except *C. c. apaporiensis* that is in Appendix I. \r\n

Please follow this link for its [CITES: legal status](#)

## Lifecycle Stages

Sexual maturity may be reached from as young as four years to as old as seven years, since less dominant specimens grow slower and sometimes don't reproduce at all. Adult males are, on average, 30 to 40% larger than adult females.

## Uses

*Caiman crocodilus* is a valuable species in the hide and pet trade industry. Their skins are the most popularly harvested product among crocodiles with over half a million traded legally each year. Since they were considered commercially inferior compared to non-ossified alligator and crocodile skins in the early 1900s, common caimans weren't hunted until the 1950s when other crocodilian populations diminished. For this reason, caimans have since remained resilient to overhunting because of harvest regulations and due to the fact that they reproduce at a relatively small size. Most countries within its native range set harvest regulations with licensing and inspection programs. Countries such as Colombia have developed captive breeding ranches to farm their hides (Ross, 1998; CSG, 2005; F. Grana, pers. comm., November 2007). Poaching for meat, and eventually for skins, is widespread throughout the Brazilian Amazonia since 1980s (Da Silveira et al., 1998; Da Silveira & Thorbjarnarson, 1999), but in only one locality is known where a *C. crocodilus* population is in danger due to subsistence hunting undertaken by indigenous people in the north of Amapa state (R. Da Silveira, pers. comm., January 2011).

## Habitat Description

*Caiman crocodilus* basically occurs in all low wetland habitats in Central and South America except along small streams in the pristine terra firme (never flooded) Amazonian forests. It commonly inhabits flooded forests, swamps, large and small rivers, lakes and canals and can occur in disturbed and polluted water bodies.\r\nIt is known to occur in any of freshwater body, natural or man-made, as long as it is warm, above 28.5°F, and deep enough to submerge itself completely. They can, on occasion, inhabit brackish waters and leave the water to warm in the sun on shores. In Puerto Rico, caimans have established in rural, suburban, and urban settings. In Manaus city, Central Amazonia, it is common and nesting along small urban polluted streams (R. Da Silveira, pers. Comm., January 2011). They are highly adaptable to new environments and readily colonize newly formed waters.

## Reproduction

Wild *C. crocodilus* populations reproduce sexually. Courtship and copulation take place in the late rainy season/early dry season. Female build a mound-nest with leaves, branches or grass. The female lays from seven to 41 eggs. The incubation period lasts from 70 to 90 days, and the female can guard and open the nest for the hatchlings to emerge. Hatchlings measure, at birth, from 17 to 25 cm long. Young associate in sibling groups or pods (multiple hatchlings from different nests) and remain close to a female that can take on maternal duties for about a year and a half.

## Nutrition

*Caiman crocodilus* are generalist and opportunistic, feeding upon a great variety of invertebrates and vertebrates. Juveniles consume mostly aquatic invertebrates such as insects, spiders, crustaceans and molluscs. Adults feed on larger prey as fish, and sometimes amphibians, reptiles, birds and mammals (Da Silveira & Magnusson, 1999; CSG, 2002; Somma, 2008). In urban habitats of the Manaus city, the diet of the species is composed mainly of *Rattus* sp. (R. Da Silveira, pers.comm., January 2011).

## General Impacts

*Caiman crocodilus* is a generalist and opportunistic predator, but due their relative small size and lack of aggressive behaviour they do not in general represent a danger for humans, pets and farm animals.

## Management Info

Physical: Within some of its native range such as in Venezuela, Guyana, and Nicaragua *C. crocodilus* wild populations are cropped, or harvested. Venezuela which operates the largest management program allows private land owners to harvest up to 15% of adult males over 180 cm each year. The program includes a strict licensing and a centralized inspection agency, but independent surveys have indicated overexploitation in some areas. Such sustainable programs require substantial survey, study, and continued monitoring. In Puerto Rico, reported caimans are captured by government officials and taken to a retention center from which they are either exported or killed (Ross, 1998; Felix Grana, pers.comm., November 2007).

## Pathway

**Principal source:** [Somma, 2008. USGS Nonindigenous Aquatic Species \(NAS\): \*Caiman crocodilus\* \(Linnaeus, 1758\);](#)

[Gulf States Marine Fisheries Commission \(GSMFC\) Non-Native Aquatic Species in the Gulf of Mexico and South Atlantic Regions: \*Caiman crocodilus\* \(Linnaeus\);](#)

[Crocodilian Biology Database, 2002. \*Caiman crocodilus\* \(LINNAEUS, 1758\) hosted by the IUCN SSC Crocodile Specialist Group](#)

[Velasco, A. & Ayarzagüena, J. 2010. Spectacled Caiman \*crocodilus\*. In S.C. Manolis and C. Stevenson \(Eds\). Crocodiles. Status Survey and Conservation Action Plan \(pp. 10-15\). Crocodile Specialist Group: Darwin.](#)

**Compiler:** National Biological Information Infrastructure (NBII), Felix A. Grana Raffucci, Technical Advisor, Puerto Rico Department of Natural & Environmental Resources & IUCN SSC Invasive Species Specialist Group (ISSG)

## Review:

Ronis Da Silveira, Departamento de Biologia, Instituto de Ciências Biológicas, Universidade Federal do Amazonas & Boris Marioni, Instituto Piagaçu, Coordinator of Caiman Project

**Publication date:** 2011-01-18

## ALIEN RANGE

**[1]** CUBA

Global Invasive Species Database (GISD) 2025. Species profile *Caiman crocodilus*. Available from: <https://www.iucngisd.org/gisd/species.php?sc=1206> [Accessed 02 July 2025]

**[5]** PUERTO RICO

Pag. 3

**[1] THAILAND**

**BIBLIOGRAPHY**

37 references found for **Caiman crocodilus**

**Management information**

- Campos, Z., Magnusson, W.E., Sanaiotti, T. & Coutinho, M. 2008. Reproductive trade-off in *Caiman crocodilus crocodilus* and *Caiman crocodilus yacare*: implications for size-related management quotas. *Herpetological Journal*, 18: 91-96.
- da Silveira, R., Gordo, M., Marcon, J. L. & Silva, J. R. 1998. Skins from Wild Spectacled Caimans Confiscated in the Amazonia. *Newsletter Crocodile Specialist Group Iucn Ssc*, Gainesville, 17(3): 7-8.
- da Silveira, R.; Magnusson, W.E. & Campos, Z. 1997. Monitoring the Distribution, Abundance and Breeding Areas of *Caiman crocodilus crocodilus* and *Melanosuchus niger* in the Anavilhas Archipelago, Central Amazonia, Brazil. *Journal of Herpetology*, 31(4): 514-520.
- da Silveira, R., Magnusson, W.E. & Thorbjarnarson, J. 2008. Factors Affecting the Number of Caimans Seen During Spotlight Surveys in the Mamirauá Reserve, Brazilian Amazonia. *Copeia*, 2008(2): 425-430.
- da Silveira, R. & Thorbjarnarson, J. 1999. Conservation Implications of Commercial Hunting of Black and Spectacled Caiman in the Mamirauá Sustainable Development Reserve, Brazil. *Biological Conservation*, 88: 103-109.

[Gulf States Marine Fisheries Commission \(GSMFC\), 2005. Non-Native Aquatic Species in the Gulf of Mexico and South Atlantic Regions: Caiman crocodilus \(Linnaeus\)](#)

**Summary:** Available from: [http://nis.gsmfc.org/nis\\_factsheet.php?toc\\_id=207](http://nis.gsmfc.org/nis_factsheet.php?toc_id=207) [Accessed 19 November 2007]

[Somma, Louis A., 2008. Caiman crocodilus. USGS Nonindigenous Aquatic Species Database, Gainesville, FL](#)

**Summary:** Available from: <http://nas.er.usgs.gov/queries/FactSheet.asp?speciesID=222> [Accessed 19 November 2007]

[Velasco, A. & Ayarzagüena, J. 2010. Spectacled Caiman crocodilus. In S.C. Manolis and C. Stevenson \(Eds\). \*Crocodiles. Status Survey and Conservation Action Plan\* \(pp. 10-15\). Crocodile Specialist Group: Darwin.](#)

**Summary:** Available from: [http://www.iucncsg.org/ph1/modules/Publications/ActionPlan3/ap2010\\_03.html](http://www.iucncsg.org/ph1/modules/Publications/ActionPlan3/ap2010_03.html) [Accessed 19 January, 2011]  
[Witmer, Gary W., Patrick W. Burke, Will C. Pitt, Michael L. Avery., 2007. Management of Invasive Vertebrates in the United States: An Overview. USDA National Wildlife Research Center Symposia Managing Vertebrate Invasive Species University of Nebraska - Lincoln Year 2007](#)

**Summary:** Available from: <http://digitalcommons.unl.edu/cgi/viewcontent.cgi?article=1055&context=nwrcinvasive> [Accessed 10 September 2008]

**General information**

Ayarzagüena, J. & Castroviejo, J. 2008. La baba (*Caiman crocodilus*) en la Estación Biológica El Frío (Estado Apure). In J. Castroviejo, J. Ayarzagüena and A. Velasco (Eds). *Contribución al Conocimiento del Género Caimán de Suramérica* (pp. 181-294). Publ. Asoc. Amigos de Doñana 18: Seville, Spain.

Behler, J.L. 1979. The Audubon Society Field Guide to North American Reptiles and Amphibians. Alfred A. Knopf, Inc. New York. 743 pp.  
Brochua, Christopher A., Ángel M. Nieves-Rivera, Jorge Vélez-Juarbe, Juan D. Daza-Vaca and Hernán Santos., 2007. Tertiary crocodylians from Puerto Rico: Evidence for Late Tertiary endemic crocodylians in the West Indies? [Crocodyliens tertiaires de Porto Rico : preuve de l'existence de crocodiliens endémiques dans les Antilles ?Cocodrilos terciarios de Puerto Rico: prueba de la existencia de cocodrilos endémico en las Antillas?] *Geobios Volume 40*, Issue 1, January–February 2007, Pages 51-59

Conant, R., and J.T. Collins. 1991. Reptiles And Amphibians. Eastern/Central North America. Houghton Mifflin Company. Boston. 450 Pp.  
[Crocodile Specialist Group 1996. Crocodylus rhombifer. In: IUCN 2007. 2007 IUCN Red List of Threatened Species.](#)

**Summary:** Available from: <http://www.iucnredlist.org/search/details.php/5670/all> [Accessed 10 September 2008]

[Crocodilian Biology Database 2002. Caiman crocodilus \(LINNAEUS, 1758\) hosted by the IUCN SSC Crocodile Specialist Group \(CSG\)](#)

**Summary:** Available from: [http://www.flmnh.ufl.edu/cnhc/csp\\_ccro.htm](http://www.flmnh.ufl.edu/cnhc/csp_ccro.htm) [Accessed 19 November 2007]  
da Silveira, R. 2011. Management of Wildlife in the Floodplain: A Critical Look at Threats, Bottlenecks, and the Future in Amazonia. In Miguel A. Pinedo-Vasquez, Mauro Ruffino, Christine J. Padoch and Eduardo S. Brondum (Eds). *The Amazon Várzea The Amazon Várzea The Decade Past and the Decade Ahead. Part 2* (pp. 137-144).

da Silveira, R. & Magnusson, W.E. 1999. Diets of Spectacled and Black Caiman in the Anavilhas Archipelago, Central Amazonia, Brazil. *Journal of Herpetology*, Estados Unidos, 33(2): 181-192.

da Silveira, R., Ramalho, E.E.; Magnusson, W.E. & Thorbjarnarson, J. 2010. Depredation by Jaguars on Caimans and Importance of Reptiles in the Diet of Jaguar. *Journal of Herpetology*, 44(3): 418-424.

Farias, I.P., da Silveira, R., de Thoisy, B., Monjelón, L.A., Thorbjarnarson, J. & Hrbek, T. 2004. Genetic diversity and population structure of Amazonian crocodylians. *Animal Conservation* 7: 265-272

[Florida Fish and Wildlife Conservation Commission., undated. Checklist of Florida's Amphibians and Reptiles.](#)

**Summary:** Available from: <http://www.fs.fed.us/outdoors/naturewatch/implementation/Wildlife/chcklst-florida%20amphibs-reptiles.pdf> [Accessed 10 September 2008]

[Forero-Medina, German; Castaño-Mora, Olga Victoria; Rodríguez-Melo, Miguel., 2006. Ecology of Caiman crocodilus fuscus on San Andres Island, Colombia: A preliminary study. Caldasia. 28\(1\). JUN 30 2006. 115-123](#)

**Summary:** Fulltext in Spanish.

Available from: <http://www.scielo.org.co/pdf/cal/v28n1/v28n1a11.pdf> [Accessed 8 September 2008]

Hertner, George, 2006. Caiman bite. *Wilderness & Environmental Medicine*. 17(4). WIN 2006. 267-270.

[ITIS \(Integrated Taxonomic Information System\), 2005. Online Database Caiman crocodilus](#)

**Summary:** An online database that provides taxonomic information, common names, synonyms and geographical jurisdiction of a species. In addition links are provided to retrieve biological records and collection information from the Global Biodiversity Information Facility (GBIF) Data Portal and bioscience articles from BioOne journals.

Available from: [http://www.itis.gov/servlet/SingleRpt/SingleRpt?search\\_topic=TSN&search\\_value=174370](http://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=174370) [Accessed 20 November 2007]

Marioni, B., da Silveira, R., Magnusson, W.E. & Thorbjarnarson, J. 2008. Feeding Behaviour of Two Sympatric Caiman Species, *Melanosuchus niger* and *Caiman crocodilus*, in the Brazilian Amazon. *Journal of Herpetology*, 42: 768-772.

[National Center for Biotechnology Information \(NCBI\) undated. Taxonomy Browser Caiman crocodilus](#)

**Summary:** Available from:

<http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=8499&lvl=3&lin=f&keep=1&srchmode=1&unlock> [Accessed 19 November 2007]

Ouboter, P.E. & Nanhoes, L.M.R. 1988. Habitat selection and migration of *Caiman crocodilus crocodilus* in a swamp and swamp-forest habitat in Northern Suriname. *Journal of Herpetology*, 2: 283-294.

[Reptiles Database, 2010. Caiman crocodilus Linnaeus, 1758](#)

**Summary:** Available from: <http://reptile-database.reptarium.cz/species.php?genus=Caiman&species=crocodilus> [Accessed September 8 2010]

Rivero, J. A. 1998. The Amphibians and Reptiles of Puerto Rico. 2nd Ed. Univ Puerto Rico Editorial. 369-371

[Ross, 1998 J.P. Ross, Crocodiles. Status survey and conservation action plan \(Second Edition\), International Union for the Conservation of Nature, Gland, Switzerland \(1998\).](#)

**Summary:** Available from: <http://www.flmnh.ufl.edu/natsci/herpetology/act-plan/plan1998a.htm#Contents> [Accessed 10 September 2008]

Schwartz, A & Henderson, R.W., 1991. Amphibians and Reptiles of the West Indies: Descriptions, distributions and natural history, University of Florida Press, Gainesville (1991).

Smith, Hobart M.& Anthony J. Kohler., 1977. A Survey of Herpetological Introductions in the United States and Canada. *Transactions of the Kansas Academy of Science* (1903-), Vol. 80, No. 1/2 (Spring - Summer, 1977), pp. 1-24 Published by: Kansas Academy

[Smithsonian National Zoological Park, 2000. Cuban Crocodile \*Crocodylus rhombifer\*. Friends of the National Zoo.](#)

**Summary:** Available from: <http://nationalzoo.si.edu/Animals/Amazonia/Facts/fact-cubancrocodile.cfm> [Accessed 10 September 2008]

[UNEP-WCMC Species Database., 2008. Caiman crocodilus \(Linnaeus, 1758\)](#)

**Summary:** Available from:

<http://www.unep-wcmc.org/isdb/Taxonomy/tax-species-result.cfm?Genus=Caiman&species=crocodilus&source=animals&tabname=distribution> [Accessed 10 September 2008]

[Vidthayanon, Chavalit., Country reports Aquatic alien species in Thailand \(Part 1\): Biodiversity](#)

**Summary:** Available from: <ftp://ftp.fao.org/docrep/fao/008/a0113e/a0113e11.pdf> [Accessed 10 September 2008]

[Williams, E. H. 1995. Parasites of caimans in Puerto Rico. Crocodile Specialist Group Newsletter \(IUCN\). 14\(4\):18-19.](#)

**Summary:** Available from: <http://www.flmnh.ufl.edu/natsci/herpetology/newsletter/news144c.htm> [Accessed 19 November 2007]

[Witmer, Gary W. & Jeffrey C. Lewis., 2001. Introduced Wildlife of Oregon and Washington. Wildlife Damage Management, Internet Center for USDA National Wildlife Research Center - Staff Publications. University of Nebraska - Lincoln](#)

**Summary:** Available from: [http://digitalcommons.unl.edu/cgi/viewcontent.cgi?article=1652&context=icwdm\\_usdanwrc](http://digitalcommons.unl.edu/cgi/viewcontent.cgi?article=1652&context=icwdm_usdanwrc) [Accessed 10 September 2008]

Ximena, Bernal, E., 2006. *Caiman crocodilus* (spectacled caiman). Predation/eavesdropper. *Herpetological Review*. 37(4). DEC 2006. 460-461.