

Physalis peruviana 简体中文 正體中文

System: Terrestrial

Kingdom	Phylum	Class	Order	Family
Plantae	Magnoliophyta	Magnoliopsida	Solanales	Solanaceae

Common name manini fuu lalahi (Niuean), aguaymanto (Spanish, Peru), Kapstachelbeere (English), coqueret du Peru (English), tomatinho-de-capucho (Portuguese), gooseberry tomato (English), erva-noiva-do-peru (Portuguese), alquequenje (Portuguese, Brazil), groselha-do-Peru (Portuguese, Brazil), manini (Niuean), alquequenje (Spanish), bate-testa (Portuguese, Brazil), capulí (Spanish), physalis (Portuguese, Brazil), uvilla (Spanish, Ecuador), tukiyandra (Fijian), botebote yadra (Fijian), poha (Hawaiian), thol thakkali (Sinhalese), topotopo (Quechua), alquequenje amarillo (Spanish), mbotembote yandra (Fijian), maulanggu (Fijian), kospeli (Fijian), tupera (Maori, Cook Islands), goundou-goundou (English), pa'ina (Hawaiian), winebusupén (English), Cape gooseberry (English), gooseberry-tomato (English), te bin (English, Kiribati), te baraki (English, Kiribati), rasabarii (Nepali), jangalii mevaa (Nepali), ishmagol (Nepali), Peruvian cherry (English), watamo (English, Nauru), ground cherry (English), Peruvian ground-cherry (English), tupere (English, Tahiti), goldenberry (English), oatamo (English, Nauru), camapú (Portuguese, Brazil), ku'usi (Tongan)

Synonym

Similar species

Summary

Physalis peruviana originates from the tropics and is cultivated in its native lands. It poses an indirect threat to agriculture when imported as it may harbour introduced plant pests.



[view this species on IUCN Red List](#)

Species Description

Physalis peruviana is a shortlived perennial to 1 m tall, semi-woody. Leaves simple, alternate, in pairs at each node, unequal in size, margins somewhat lobed or not. Flowers solitary in leaf axils, bell-shaped, yellow with purplish brown spots at the base of the corolla. Berries pale yellow, drying to pale brown, aromatic, covered by papery calyx. Seeds numerous (Motooka *et al.* 2003).
\\n

Soft-wooded, short-lived shrubs up to ca. 1 m tall, straggly with age, all parts densely pubescent with erect, simple or glandular hairs up to 1 mm long. Leaves simple, alternate, usually geminate, 1 larger than the other, ovate-acuminate, often 6 cm long, 4 cm wide, margins entire or rarely with a few blunt lobes, apex acuminate, base cordate, petioles 2 to 3 cm long. Flowers perfect, actinomorphic, solitary in the leaf axils, pedicellate; calyx connate in lower, 5-lobed, veins often prominent, the lobes acuminate-triangular, ca. 1 cm long, distinct at apex; corolla yellow with well-defined purplish brown spots at base, 15 to 20 mm in diameter, the limb rotate or shallowly 10-lobed, the tube swollen into shallow nectary pouches between the filaments, densely pubescent with pale yellowish dendritic hairs below the spots and around the nectaries; style 5 to 7 mm long. Berries pale yellow, drying pale brown, aromatic, succulent, globose, 1.5 to 2 cm in diameter, enclosed in the inflated calyx 3 to 3.5 cm long. Seeds numerous, pale brown, discoid, 1.75 to 2 mm long, minutely shallowly reticulate, embryo curved, endosperm present (Wagner *et al.* 1999, in PIER 2002).
\\n

Seeds creamy white to yellowish and very small (less than 2 mm in diameter); ovoid, compressed; testa slightly pitted (PIER Undated). \\n

Notes

The scientific name *Physalis peruviana* was derived from the Greek *physa*, bladder, for the calyx covering the fruit and *peruviana* meaning of Peru (Motooka *et al.* 2003).\\n

Uses

Physalis peruviana is used as an ornamental plant; consumed (fruit); berries used for making jams; used in traditional medicine (USDA-ARS 2003; Motooka *et al.* 2003).

Habitat Description

Physalis peruviana grows well in the tropics (Bailey 1949, in USDA 1997). It may be found in mesic to wet forests, subalpine woodland and disturbed sites on mountain slopes at altitudes of 450 to 2020 meters (Wagner *et al.* 1999, in PIER 2002; Motooka *et al.* 2003). In Fiji it occurs at elevations from near sea level to 900 meters; it is also found in gardens and in forests along trails and streams, in clearings and in cultivated areas (Smith 1991, in PIER 2002). It is a common weed in some plantations in Niue (Sykes 1970, in PIER 2002). It is frequently found in Tahiti in cool valleys to an altitude of 800 meters (Welsh 1998, in PIER 2002). It is an occasional plantation weed in Tonga (Yuncker 1959, in PIER 2002).

Reproduction

Fruit/seed; seed produced in a fruit capsule

General Impacts

The Global Compendium on Weeds lists *Physalis peruviana* as an agricultural weed, cultivation escape and environmental weed (GCW 2007). *P. peruviana* poses an indirect threat to US agriculture when imported as it may harbour introduced plant pests, including a wide range of arthropod (insect) pests and plant pathogenic fungi, viruses and bacteria (USDA 1997).

Management Info

Chemical: *Physalis peruviana* is probably susceptible to hormone-type herbicides, especially when young, and probably to tebuthiuron (Motooka *et al.* 2003).

Principal source:

Compiler: IUCN SSC Invasive Species Specialist Group (ISSG) with support from the EU-funded South Atlantic Invasive Species project, coordinated by the Royal Society for the Protection of Birds (RSPB)

Review:

Publication date: 2010-08-16

ALIEN RANGE

[1] AFRICA	[1] ASIA
[4] AUSTRALIA	[1] AUSTRIA
[1] BERMUDA	[1] CANADA
[3] COOK ISLANDS	[1] DENMARK
[1] ECUADOR	[3] FIJI
[5] FRENCH POLYNESIA	[1] GUAM
[1] HUNGARY	[1] INDIA
[1] INDONESIA	[2] JAPAN
[1] KENYA	[1] KIRIBATI
[1] NAURU	[4] NEW CALEDONIA
[3] NEW ZEALAND	[1] NIUE
[1] NORFOLK ISLAND	[2] PALAU
[1] PITCAIRN	[1] PORTUGAL
[1] REUNION	[3] SAINT HELENA
[1] SEYCHELLES	[1] TAIWAN
[3] TONGA	[1] TUVALU
[1] UNITED KINGDOM	[12] UNITED STATES
[1] WEST INDIES	[1] ZIMBABWE

BIBLIOGRAPHY

17 references found for ***Physalis peruviana***

Management information

[Global Compendium of Weeds \(GCW\). 2007. *Physalis peruviana* \(Solanaceae\)](#)

Summary:

Available from: http://www.hear.org/gcw/species/physalis_peruviana/ [Accessed 25 October 2009]

[IUCN/SSC Invasive Species Specialist Group \(ISSG\)., 2010. A Compilation of Information Sources for Conservation Managers.](#)

Summary: This compilation of information sources can be sorted on keywords for example: Baits & Lures, Non Target Species, Eradication, Monitoring, Risk Assessment, Weeds, Herbicides etc. This compilation is at present in Excel format, this will be web-enabled as a searchable database shortly. This version of the database has been developed by the IUCN SSC ISSG as part of an Overseas Territories Environmental Programme funded project XOT603 in partnership with the Cayman Islands Government - Department of Environment. The compilation is a work under progress, the ISSG will manage, maintain and enhance the database with current and newly published information, reports, journal articles etc.

[Motooka, P. et al. 2003. Weeds of Hawai'i's Pastures and Natural Areas; An Identification and Management Guide. College of Tropical Agriculture and Human Resources, University of Hawaii: Manoa.](#)

Summary: Available from: http://www.ctahr.hawaii.edu/forestry/Data/WeedsHI/W_Physalis_peruviana.pdf [Accessed 25 October 2009]
[Pacific Island Ecosystems at Risk \(PIER\). 2008. Images: *Physalis peruviana* L., Solanaceae](#)

Summary:

Available from: http://www.hear.org/pier/imagepages/thumbnails/physalis_peruviana.htm [Accessed 25 October 2009]

[Tristan Island Government. 2006. Tristan da Cunha: Biodiversity Action Plan \(2006-2010\)](#)

Summary: Available from: http://www.rspb.org.uk/Images/TristanBiodiversityActionPlan2_tcm9-180968.pdf [Accessed 25 October 2009]
[U.S. Department of Agriculture \(USDA\). 1997. Importation of Cape Gooseberry Fruit, *Physalis peruviana*, from Colombia into the United States: Qualitative, Pathway-Initiated Pest Risk Assessment.](#)

Summary:

Available from: [https://web01.aphis.usda.gov/oxygen_fod/fb_md_ppq.nsf/d259f66c6afbd45e852568a90027bcad/cf02d9a32e4823d4852568f60048c304/\\$FILE/0023.pdf](https://web01.aphis.usda.gov/oxygen_fod/fb_md_ppq.nsf/d259f66c6afbd45e852568a90027bcad/cf02d9a32e4823d4852568f60048c304/$FILE/0023.pdf) [Accessed 25 October 2009]

[Varnham, K. 2006. Non-native species in UK Overseas Territories: a review. JNCC Report 372. Peterborough: United Kingdom.](#)

Summary: This database compiles information on alien species from British Overseas Territories.

Available from: <http://www.jncc.gov.uk/page-3660> [Accessed 10 November 2009]

General information

Global Invasive Species Database (GISD) 2026. Species profile *Physalis peruviana*. Available from:

<https://www.iucngisd.org/gisd/species.php?sc=1461> [Accessed 05 February 2026]

[Ascension Island Conservation Centre. Undated. Plants. Georgetown, Ascension Island, South Atlantic Ocean.](#)

Summary: Available from: <http://www.ascensionconservation.org.ac/pdf/6-E-plants-of-Green-Mountain.pdf> [Accessed 25 October 2009]
Dean, W.R.J., S.J. Milton, P.G. Ryan and C.L. Moloney. 1994. The role of disturbance in the establishment of indigenous and alien plants at Inaccessible and Nightingale Islands in the South Atlantic Ocean, *Plant Ecology* 113 (1)

[Duffey, Eric. 1964. The Terrestrial Ecology of Ascension Island, The Journal of Applied Ecology 1 \(2\): 219-251.](#)

Summary: Images

Available from: http://www.seaturtle.org/PDF/Duffey_1964_JAppEcol.pdf [Accessed 25 October 2009]

Fay, Michael F., Vanessa E. Thomas and Sandra Knapp. 2007. 602. *Mellissia begonifolia* Solanaceae. The Board of Trustees of the Royal Botanic Gardens, Kew 2007. Published by Blackwell Publishing Ltd

[Gray, Alan, Tara Pelembe and Stedson Stroud. 2005. The conservation of the endemic vascular flora of Ascension Island and threats from alien species, *Oryx* 39 \(4\)](#)

[Hawaiian Ecosystems at Risk project \(HEAR\). 2008. Plants of Hawaii: Solanaceae: *Physalis peruviana* \(Poha, Cape gooseberry\)](#)

Summary: Images

Available from: <http://www.hear.org/starr/plants/images/species/?q=physalis+peruviana> [Accessed 25 October 2009]

[Medeiros, A.C., L.L. Loope and C.G. Chimera. 1998. Flowering Plants and Gymnosperms of Haleakala National Park \[Technical Report 120\]. Department of Botany, Pacific Cooperative Studies Unit University of Hawaii: Manoa.](#)

Summary: Images

Available from: <http://www.botany.hawaii.edu/faculty/duffy/techr/120.pdf> [Accessed 25 October 2009]

[Pacific Island Ecosystems at Risk \(PIER\). 2002. *Physalis peruviana* L., Solanaceae](#)

Summary: Available from: http://www.hear.org/pier/species/physalis_peruviana.htm [Accessed 25 October 2009]

[United States Department of Agriculture - Agricultural Research Service \(USDA-ARS\). 2003. Taxon: *Physalis peruviana* L.. National Genetic Resources Program. Germplasm Resources Information Network - \(GRIN\) \[Online Database\]. National Germplasm Resources Laboratory, Beltsville, Maryland.](#)

Summary: Images

Available from: http://www.ars-grin.gov/cgi-bin/npgs/html/tax_search.pl?Physalis%20peruviana [Accessed 25 October 2009]

[United States Department of Agriculture - Natural Resources Conservation Service \(USDA-NRCS\). 2008. *Physalis peruviana* L. Peruvian groundcherry. The PLANTS Database. National Plant Data Center, Baton Rouge, LA 70874-4490 USA.](#)

Summary: Images

Available from: <http://plants.usda.gov/java/profile?symbol=PHPE4> [Accessed 25 October 2009]