Pitangus sulphuratus **System:** Terrestrial

Kingdom	Phylum	Class	Order	Family
Animalia	Chordata	Aves	Passeriformes	Tyrannidae

Common name

Bentevi (Portuguese), Tyran kiskidi (French), Tyran sulfureux (French), Tyran quiquivi (French), Tyran quesquildit (French), Moucherolle masqué (French), Bentevi (German), Grote kiskadie (Dutch), Naamioväijy (Finnish), Bentewi wielki (Polish), Kibaraootairanchou (Japanese), Pitango solforato (Italian), Bem-te-vi (Portuguese), Storkiskadi (Norwegian), Större kiskadi (Swedish), Bem-te-vi-verdadeiro (Portuguese), Bem-te-viu (Portuguese), Bem-te-vi-de-coroa (Portuguese), Triste-vida (Portuguese), Bentevi-de-coroa (Portuguese), Bem-te-vi-carrapateiro (Portuguese), Pitanguá (Portuguese), Pitauã (Portuguese), Pituã (Portuguese), Siririca (Portuguese), Bentevi-verdadeiro (Portuguese), Bem-te-vi-de-cabeça-rajada (Portuguese), Lord Derby's flycatcher (English), Kiskadee flycatcher (English), Derby flycatcher (English), Greater kiskaee (English), Great kiskadee (English), Pitogue (English), Luis grande (Spanish, Mexico), Luis bienteveo (Spanish, Mexico), Bienteveo grande (Spanish, Costa Rica), Cristofué (Spanish, Honduras), Pitogüé (Spanish, Paraguay), Tyran bentevi (Slovak), Postriežkár bentevi (Slovak), Güis común (Spanish, Nicuragua), Pitangua (English), Schwefelmaskentyrann (German), Kiskadie (Danish), Benteveo (Spanish, Argentinia, Bolivia, Chile), Tyran bentevi (Czech), Great Kiskadee (English), Benteví (Portuguese), Bichofué (Spanish, Columbia), Kiskadì maggiore (Italian), Benteveo común (Spanish, Argentinia, Uruguay)

Synonym

Lanius sulphuratus
Lanius sulphuratus ,Linnaeus, 1766

Similar species

Summary

view this species on IUCN Red List

Notes

While the genus *Pitangus* is monotypic, there are a number of described *P. sulphuratus* subspecies: *P. sulphuratus argentinus* (Todd, 1952), *P. sulphuratus bolivianus* (Lafresnaye, 1852), *P. sulphuratus caucensis* (Chapman, 1914), *P. sulphuratus derbianus* (Kaup, 1852), *P. sulphuratus guatimalensis* (Lafresnaye, 1852), *P. sulphuratus maximiliani* (Cabanis & Heine, 1859), *P. sulphuratus rufipennis* (Lafresnaye, 1851), *P. sulphuratus sulphuratus* (Linnaeus, 1766), *P. sulphuratus texanus* (Van Rossem, 1940),*P. sulphuratus trinitatis* (Hellmayr, 1906) (ITIS, 2010).

Uses

200 individuals of *Pitangus sulphuratus* were intentionally introduced to Bermuda from Trinidad in 1957 as biological control agents for the introduced anole lizards (particularly the Jamaican anole (*Norops grahami*) (Cheesman & Clubbe, 2007; Davenport *et al.*, 2008) and insect pests (Griffith *et al.*, unpublished data). However, this biocontrol attempt was a failure, with *P. sulphuratus* preferring to feed on various species of native fauna instead.

Habitat Description
Preferred habitats of *Pitangus sulphuratus* include around rivers, streams, and lakes bordered with dense vegetation as well as open country and parks (Field Guide to Birds of North America, 2007). *P. sulphuratus* has also been observed in arable land, pastureland, heavily degraded forest, moist lowland and montane forests, moist shrubland, dry grassland, mangrove habitats (Birdlife International, 2009); and beaches (Latino & Beltzer, 1999).

Reproduction
Pitangus sulphuratus forms monogamous relationships. It breeds in open woodland with some tall trees, including cultivated areas and around human habitation. Two to five creamy white eggs spotted with brown and lavender are laid in a bulky, domed nest with a side entrance made of grass, weeds, bark strips, moss, and other plant fibers, and built in trees, bush and telephone poles 6 to 50 feet above the ground. Incubation ranges from 13 to 15 days and is carried out by the female. (Field Guide to Birds of North America, 2007).

Nutrition
Pitangus sulphuratus is a generalist feeder, with a stomach content analysis on 83 *P. Sulphuratus* stomachs revealing that insects had the highest relative importance index score of 5304, followed by seed and fruit (4606), crustaceans (170) and fishes (154) (Latino & Beltzer, 1999). Feeding activity was the highest between 11 am and 4 pm (Latino & Beltzer, 1999). Although they are physically unable to probe in the dirt for insect larvae and other invertebrates such as earthworms and snails, kleptoparasitic behaviour has been observed, with instances of *P. sulphuratus* stealing these food items from other birds being recorded (Llambıas *et al.*, 2001). *P. sulphuratus* is also known to feed on frogs, small lizards, baby birds, mice (Field Guide to Birds of North America, 2007), and bats (Fischer *et al.*, 2010). In urban envronments, the ability that *P. sulphuratus* has for identifying food items absent from more natural habitats (such as human food scraps and animal food pellets) and for exploiting resources of unpredictable spatial and temporal distributions confers the species a dietary flexibility that probably contributes to its efficiency in colonizing urban habitats (Argel-De-Oliveira *et al.*, 1998).

General Impacts
In its introduced range, in Bermuda, *Pitangus sulphuratus* predates on and competes with various native species, playing a significant role in the population declines of insect, bird and reptile species (Cheesman & Clubbe, 2007; Davenport *et al.*, 2008; Forbes, 2010).
P. sulphuratus disperses seeds of invasive species through native forest, contributing to the spread of invasive species and the decline of native plant species (The Encyclopedia of Earth, 2006). As a generalist feeder, it is reported as an agricultural pest in its native range (Rodriguez *et al.*, 2004).

Management Info
Physical control: *Pitangus sulphuratus* populations on Nonsuch Island , Bermuda are controlled by shooting (Varnham, 2006), however the effectiveness of this is uncertain with Griffith *et al.* (unpublished data) stating that reforestation has allowed *P. Sulphuratus* to persist on the island in spite of control efforts. Physical control methods for *P. sulphuratus* and other bird pests on vineyards in Uruguay include use of firearms, visual repellants (flags and scarecrows), and acoustic repellents (fireworks, propane cannons and distress calls) (Rodriguez *et al.*, 2004).

Chemical control: Chemical control methods for *P. sulphuratus* and other bird pests on vineyards in Uruguay include use of toxic baits (Carbofuran) and repellents (methiocarb) (Rodriguez *et al.*, 2004).

Other: *Pitangus sulphuratus* has a natural aversion to patterns resembling those of coral snake species (*Micrurus spp.* and *Micruroides spp.*) (Brodie III & Janzen, 1995).

Principal source:

Compiler: IUCN SSC Invasive Species Specialist Group (ISSG) with support from the Overseas Territories Environmental Programme (OTEP) project XOT603, a joint project with the Cayman Islands Government - Department of Environment

Review: Under expert review

Pubblication date: 2010-06-02

ALIEN RANGE

[2] BERMUDA

Red List assessed species 6: CR = 1; LC = 5;

Dumetella carolinensis bermudianus **LC**
Eumeces longirostris **CR**
Phyllomedusa distincta **LC**
Physalaemus albonotatus **LC**
Scinax nasicus **LC**
Sialia sialis **LC**

BIBLIOGRAPHY

49 references found for **Pitangus sulphuratus**

Managment information

IUCN/SSC Invasive Species Specialist Group (ISSG)., 2010. A Compilation of Information Sources for Conservation Managers.
Summary: This compilation of information sources can be sorted on keywords for example: Baits & Lures, Non Target Species, Eradication, Monitoring, Risk Assessment, Weeds, Herbicides etc. This compilation is at present in Excel format, this will be web-enabled as a searchable database shortly. This version of the database has been developed by the IUCN SSC ISSG as part of an Overseas Territories Environmental Programme funded project XOT603 in partnership with the Cayman Islands Government - Department of Environment. The compilation is a work under progress, the ISSG will manage, maintain and enhance the database with current and newly published information, reports, journal articles etc.

Rodriguez, E.N., Tiscornia, G., & Tobin, M.E. 2004. Bird depredations in Uruguyan vineyards. Wildlife Damage Management, Internet Center for USDA National Wildlife Research Center - Staff Publications. University of Nebraska: Lincoln.

General information

Argel-De-Oliveira, Maria Martha; Curi, Nancy A.; Passerini, Tatiana, 1998. Feeding of a fledgling great Kiskadee, *Pitangus sulphuratus* (Linnaeus) (Passeriformes, Tyrannidae) in urban environment. Revista Brasileira de Zoologia. 15(4). Dec., 1998. 1103-1109.

AviBase, 2010. *Pitangus sulphuratus*.
Summary: Available from: http://avibase.bsc-eoc.org/species.jsp?lang=EN&avibaseid=9C883C7BCFAEB70D [Accessed 20 April 2010]

Avila, Robson W., 2005. *Scinax nasicus* (Lesser Snouted treefrog). Predation. Herpetological Review. 36(3). SEP 2005. 308.

Bacon, Jamie P.; Gray, Jennifer A.; Kitson, Lisa, 2006. Status and conservation of the reptiles and amphibians of the Bermuda islands. Applied Herpetology, Volume 3, Number 4, 2006 , pp. 323-344(22)

Bennett, Fred D. and I. W. Hughes, 1959. Biological Control of Insect Pests in Bermuda. Bulletin of Entomological Research (1959), 50:423-436 Cambridge University Press

BirdLife International 2009. *Pitangus sulphuratus*. In: IUCN 2010. IUCN Red List of Threatened Species.
Summary: Available from: http://www.iucnredlist.org/apps/redlist/details/145369/0/full [Accessed 18 April 2010]

BirdLife International 2009. Species factsheet: *Pitangus sulphuratus*.
Summary: Available from: http://www.birdlife.org/datazone/species/index.html?action=SpcHTMDetails.asp&sid=4438&m=0 [Accessed 18 April 2010]

Birkenholz, Dale E. and Ted T. Allen, 1962. Possible Change in Status of Brewer s Blackbird in Florida. The Auk, Vol. 79, No. 1 (Jan., 1962), pp. 111-112

Brodie III, E.D., & Janzen, F.J. 1995. Experimental studies of Coral Snake mimicry: generalized avoidance of ringed snake patterns by free-ranging avian predators. Functional Ecology, 9 (2), 186-190.

Castanho, Luciano Mendes, 2001. *Phyllomedusa distincta* (leaf frog) Herpetological Review. 32(2). June, 2001. 103.

Cheesman, O.D. & Clubbe, C. 2007. Dealing with Alien Invasive Species ♦ Introduction, Overview and Conclusions. pp 193-200 in Biodiversity That Matters: a conference on conservation in UK Overseas Territories and other small island communities, Jersey 6th to 12th October 2006 (ed. M. Pienkowski). UK Overseas Territories Conservation Forum, www.ukotcf.org
Summary: Available from: http://www.ukotcf.org/pdf/JerseyConf/topic6e.pdf [Accessed 18 April 2010]

Crowell, Kenneth L., 1962. Reduced Interspecific Competition among the Birds of Bermuda. Ecology, Vol. 43, No. 1 (Jan., 1962), pp. 75-88

Crowell, K.L. and Crowell, M.R. 1976. Bermuda s abundant, beleagured birds. Nat. History 85, 48-56.

Crump Martha L. and Marcos Vaira, 1991. Vulnerability of *Pleurodema borelli* Tadpoles to an Avian Predator: Effect of Body Size and Density. Herpetologica, Vol. 47, No. 3 (Sep., 1991), pp. 316-321

Davenport, John; Jeremy Hills; Anne Glasspool & Jack Ward, 2008. Threats to the Critically Endangered endemic Bermudian skink *Eumeces longirostris*. Oryx Volume 35 Issue 4, Pages 332 - 339

D Heursel, A. And C. F. B. Haddad, 1999. Unpalatability of *Hyla semilineata* tadpoles (Anura) to captive and free-ranging vertebrate predators. Ethology Ecology & Evolution 11: 339-348, 1999
Summary: Available from: http://fupress.net/index.php/eee/article/viewFile/851/797 [Accessed 18 April 2010]
Dobson, Andrew, 1999. Conservation Measures for the Eastern Bluebird in Bermuda. Bermuda Audubon Society - Bird Conservation
Summary: Available from: http://www.audubon.bm/Bird%20Conservation.htm [Accessed 18 April 2010]
Dobson, Andrew & Wardle, Carolyn, 2005. Invasive Species Working Group. Journal of Caribbean Ornithology Vol 18, Pages 107-108
Summary: Available from: http://www.bu.edu/scscb/working_groups/resources/invasives-report-jco18-2005.pdf [Accessed 18 April 2010]
Ellison, Joanna C., 1996. Pollen Evidence of Late Holocene Mangrove Development in Bermuda. Global Ecology and Biogeography Letters, Vol. 5, No. 6 (Nov., 1996), pp. 315-326
Felipe Toledo, Luis; Woehl, Germano Jr; Woehl, Elza Nishimura; Prado, Cynthia P. A., 2005. *Scinax nasicus*, *Hyla albomarginata*, *Hyla bischoffi* and *Phyllomedusa distincta* (Tree frogs): Avian predation. Herpetological Bulletin.(92). SUM 2005. 31-32.
Field Guide to Birds of North America. 2007. Great Kiskadee.
Summary: Available from http://identify.whatbird.com/obj/852/identification/Great_Kiskadee.aspx [Accessed 20 April 2010]
Fischer, Erich; Munin, Roberto L.; Longo, Jose M.; Fischer, Wagner; de Souza, Paulo R., 2010. Predation on bats by Great Kiskadees. Journal of Field Ornithology. 81(1). MAR 2010. 17-20
Fonaroff, L. Shcuyler, 1974. Urbanization, birds and ecological change in northwestern Trinidad. Biological Conservation, Volume 6, Issue 4, Pages 258-262
Forbes, K.A. (2010). Bermuda s Fauna. Bermuda Online
Summary: Available from http://www.bermuda-online.org/fauna.htm [Accessed 20 April 2010].
Global Biodiversity Information Facility (GBIF), 2010. Subspecies: *Pitangus sulphuratus* cancensis
Summary: Available from: http://data.gbif.org/species/22179292/ [Accessed 15 June 2010]
Government of Bermuda, Ministry of Environment and Sports, Department of Conservation Services, 2009. Kiskadee (*Pitangus sulphuratus*)
Summary: Available from: http://www.conservation.bm/kiskadee/ [Accessed 18 April 2010]
Griffith, H., Wingate, D.B., & Robinson, D.L. (unpublished data).
Summary: Available from http://www.docstoc.com/docs/31994359/Introduced-Birds-as-Probable-Agents-of-Population-Changes-in-the [Accessed 20 April 2010]
Howarth, G. Francis, 1991. Environmental Impacts of Classical Biological Control. Annu. Rev. Entomol. 1991. 36:485-509
Integrated Taxonomic Information System (ITIS), 2009. *Pitangus sulphuratus* (Linnaeus, 1766)
Summary: Available from: http://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=178301 [Accessed 20 April 2010]
IUCN 2010. IUCN Red List of Threatened Species. *Phyllomedusa distinca*
Summary: Available from http://www.iucnredlist.org/apps/redlist/details/55849/0 [Accessed 23 April 2010]
Lago-Paiva, Celso, 1996. Cavity Nesting by Great Kiskadees (*Pitangus sulphuratus*): Adaptation or Expression of Ancestral Behavior? Short Communication and Commentaries. The Auk 113(4):953-955, 1996
Summary: Available from: http://elibrary.unm.edu/sora/Auk/v113n04/p0953-p0955.pdf [Accessed 18 April 2010]
Latino, Santiago; Beltzer, Adolfo, 1999. Trophic ecology of great-kiskadee *Pitangus sulphuratus* (Aves: Tyrannidae) in the Parana River Foodplain. Argentina. ORSIS.(14). 1999. 69-78.
Summary: Available from: http://ddd.uab.cat/pub/orsis/02134039v14p69.pdf [Accessed 18 April 2010]
Llambias, Paulo E. and Valentina Ferretti, 2003. Parental Care in the Great Kiskadee. The Wilson Bulletin, Vol. 115, No. 2 (Jun., 2003), pp. 214-216
Llambias, Paulo E.; Ferretti, Valentina; Rodriguez, Pablo S., 2001. Kleptoparasitism in the Great Kiskadee. Wilson Bulletin. 113(1). March, 2001. 116-117.
Lockwood, Julie L. and Michael P. Moulton, 1994. Ecomorphological pattern in Bermuda birds: the influence of competition and implications for nature preserves. Evolutionary Ecology, 1994, 8, 53-60
Mathys, Blake A.; Lockwood, Julie L., 2009. Rapid evolution of great kiskadees on Bermuda: an assessment of the ability of the island rule to predict the direction of contemporary evolution in exotic vertebrates. Journal of Biogeography. 36(12). DEC 2009. 2204-2211
Moralez-Silva, Emmanuel; Monteiro-Filho, Emygdio L. A., 2008. Kleptoparasitism by Great Kiskadee on Little Blue Heron. Waterbirds. 31(4). DEC 2008. 666-667.
Moulton, Michael P. and James G. Sanderson, 1997. Predicting the Fates of Passeriform Introductions on Oceanic Islands. Conservation Biology, Vol. 11, No. 2 (Apr., 1997), pp. 552-558
Rodrigues, Domingos J.; Arruda, Rafael, 2006. *Scinax fuscovarius* (snouted treefrog), *Scinax cf. ruber*, and *Physalaemus albonotatus* (Menwig frog). Predation. Herpetological Review. 37(2). JUN 2006. 212.
Simberloff, Daniel; Don C. Schmitz; Tom C. B. 1997. Strangers in paradise: impact and management of nonindigenous species in Florida
Summary: In Bermuda, nest boxes for eastern bluebirds (Sialia sialis) are used as perches by the non-indigenous great Kiskadee, which prevents bluebirds from nesting. (Samuel 1975 in
Smith, M. Susan, 1978. Predatory behaviour of young great kiskadees (*Pitangus sulphuratus*). Animal Behaviour Volume 26, Part 4, November 1978, Pages 988-995
Smith, Paul 2006. Interspecific Aggression in Nesting Great Kiskadees (*Pitangus sulphuratus*) in Paraguay. Bolet&#65533;n SAO Vol. XVI (No. 01) - Jul. 2006
Summary: Available from: http://www.sao.org.co/publicaciones/boletinsao/09SmithPitangus.pdf [Accessed 18 April 2010]
Stevenson, Henry M., 1972. The Recent History of Bachman s Warbler. The Wilson Bulletin, Vol. 84, No. 3 (Sep., 1972), pp. 344-347
The Birds of North America, undated. Distribution of the Great Kiskadee in North and Middle America.
Summary: Available from: http://bna.birds.cornell.edu/bna.html/species/622/galleries/figures/figure-1/image_popup_view [Accessed 18 April 2010]
The Encyclopedia of Earth (EOL), 2006. Bermuda subtropical conifer forests. World Wildlife Fund (WWF) This article has been reviewed and approved by the following Topic Editor: Mark McGinley
Summary: Available from: http://www.eoearth.org/article/Bermuda_subtropical_conifer_forests [Accessed 18 April 2010]

UniProt Taxonomy, 2010. Species *Pitangus sulphuratus* (Great kiskadee)
Wingate, David B., 1965. Terrestrial Herpetofauna of Bermuda. Herpetologica, Vol. 21, No. 3 (Sep. 24, 1965), pp. 202-218