

Norops grahami

System: Terrestrial

Kingdom	Phylum	Class	Order	Family
Animalia	Chordata	Reptilia	Squamata	Polychrotidae

Common name Graham's anole (English), common lizard (English), Jamaican anole (English)

Synonym *Anolis grahami* , Gray, 1845
Anolis iodurus , Gosse, 1850
Anolis punctatissimus , Hallowell, 1856
Anolis heterolepis , Hallowell, 1856

Similar species

Summary The Jamaican anole *Norops grahami* was introduced to Bermuda from Jamaica in 1905 to reduce populations of the fruit fly ([Ceratitis capitata](#)). In 1958 it was observed to predate heavily on beneficial insect species brought in to control introduced scale insects, subsequently resulting in the introduction of the great kiskadee ([Pitangus sulphuratus](#)), now a serious threat in itself. More recently *N. grahami* has been observed to predate on, and compete with juveniles of the 'Critically Endangered (CR)' Bermudian rock lizard ([Eumeces longirostris](#)).



[view this species on IUCN Red List](#)

Species Description

Norops grahami has a mean snout to vent length of 68.9 mm for males (Losos, 1996).

Notes

Subspecies *Norops grahami grahami* is reported from western Jamaica and Cabarita Island off Port Maria; and *N. g. aquarium* from Portland and St. Thomas parishes, Jamaica (Reptiles Database, 2010).

N. grahami is the most widespread and common of three introduced anole lizards present on Bermuda; the others are the Barbuda Bank tree anole (see [Anolis leachii](#)) and the Barbados anole ([Anolis extremus](#)) (Wingate, 1965). The effects of these lizards, particularly *N. grahami* led to the introduction of the great kiskadee ([Pitangus sulphuratus](#)) as a biocontrol agent in 1957. However, this biocontrol attempt was a failure; *P. sulphuratus* has been implicated in the population declines of native insect, bird and reptile species on Bermuda (Cheesman & Clubbe, 2007; Davenport et al., 2008).

Uses

Norops grahami was first introduced to Bermuda as a biocontrol agent to reduce populations of the fruit fly ([Ceratitis capitata](#)) in 1905 (Bennett & Hughes, 1959; Wingate, 1965).

Habitat Description

Although primarily aboreal, *Norops grahami* occupies a very wide range of habitats on Bermuda. It is observed commonly on trunks and branches of trees, on houses, stone walls, fences, piles of timber or rubble, in the taller weeds of open fields, and on open ground up to 25 feet from cover (Wingate, 1965). This species remains active all year on Bermuda although on windy overcast days it generally takes shelter near the ground in dense shrubbery, grasses, or in rubble walls. It commonly takes refuge from danger by running down into matted ground cover and crevices of walls, but will also escape upwards in larger shrubs and trees (Wingate, 1965).

Nutrition

The diet of *Norops grahami* was found to consist almost entirely of minute foliage insects (Simmonds, 1958). In order of abundance these included: Hymenoptera, especially *Brachymyrmex*, *Pheidole*, *Iridomyrmex* (Formicidae), *Aphytes* (Eulophidae) and *Chalcids* (Chalcidae). Homoptera, especially *Aphis* (Aphidae), *Pulvinaria* (Coccidae) and *Psocidae*. Diptera, various species, especially *Hippelates*. Coleoptera, various small species, especially *Ayza*, *Cybocephalus*, *Lindorus* (Coccinellidae). Lepidoptera, various species, adults and larvae (Simmonds, 1958). There was little seasonal variation in either quantity or variety of food taken, but individuals exhibited great variation in the composition of their diet; this was thought to be due to local variation in insect populations and possible habituation of some individuals to eating one particular species (Simmonds, 1958). In addition to insects, *N. grahami* has also been observed to feed on juveniles of native skinks and the eggs of native birds on Bermuda (see General Impacts).

General Impacts

Noted to feed mainly on insects, *Norops grahami* and other introduced anole lizards were observed feeding heavily on insect biological control agents that were brought in to control introduced scale insect (Simmonds, 1958); this is thought to be partly responsible for the failure of the Coccinellid *Cryptolaemus montrouzieri* to establish itself (Bennett & Hughes, 1958).

N. grahami has also been observed to prey on juveniles of the 'Critically Endangered (CR)' Bermudian rock lizard (*Eumeces longirostris*) (Griffith & Wingate, 1994; in Bacon *et al.*, 2006) and predate on the eggs of the native, eastern blue bird (*Sialia sialis*) (Thomas, 2004; in Bacon *et al.*, 2006). Although primarily aboreal, juvenile *N. grahami* can be observed foraging on the ground in skink habitats, and therefore almost certainly competing directly at times with hatchling and juvenile *E. longirostris* (Edgar, pers. comm.; in Bacon *et al.*, 2006).

Management Info

Biological control: The great kiskadee (*Pitangus sulphuratus*) was introduced as a biocontrol agent for introduced anole lizards including the brown anole (*Norops sagrei*) in 1957. However, this biocontrol attempt was a failure, with *P. sulphuratus* playing a significant role in the population declines of native insect, bird and reptile species on Bermuda (Cheesman & Clubbe, 2007; Davenport *et al.*, 2008).

Pathway

Norops grahami was introduced to Bermuda to control populations of the fruit fly *Ceratitis capitata* in 1905 (Wingate, 1965) as they were believed to be responsible for suppressing various fruit fly populations on Jamaica (Bennett & Hughes, 1958).

Principal source:

Compiler: IUCN SSC Invasive Species Specialist Group (ISSG) with support from the Overseas Territories Environmental Programme (OTEP) project XOT603, a joint project with the Cayman Islands Government - Department of Environment

Review:

Publication date: 2010-06-08

ALIEN RANGE

[1] BERMUDA

BIBLIOGRAPHY

14 references found for *Norops grahami*

Management information

Horn, Scott; Hanula, James L. 2006. Burlap bands as a sampling technique for green anoles (*Anolis carolinensis*) and other reptiles commonly found on tree boles. *Herpetological Review*. 37(4). DEC 2006. 427-428

Summary: Available from: http://www.srs.fs.usda.gov/pubs/ja/ja_horn011.pdf [Accessed 2 July 2010]

[IUCN/SSC Invasive Species Specialist Group \(ISSG\), 2010. A Compilation of Information Sources for Conservation Managers.](#)

Summary: This compilation of information sources can be sorted on keywords for example: Baits & Lures, Non Target Species, Eradication, Monitoring, Risk Assessment, Weeds, Herbicides etc. This compilation is at present in Excel format, this will be web-enabled as a searchable database shortly. This version of the database has been developed by the IUCN SSC ISSG as part of an Overseas Territories Environmental Programme funded project XOT603 in partnership with the Cayman Islands Government - Department of Environment. The compilation is a work under progress, the ISSG will manage, maintain and enhance the database with current and newly published information, reports, journal articles etc.

General information

Bacon, Jamie P., Jennifer A. Gray; Lisa Kitson, 2006. Status and conservation of the reptiles and amphibians of the Bermuda islands. *Applied Herpetology*. 3: 323-344

Bennett, Fred B. & I. W. Hughes, 1959. Biological Control of Insect Pests in Bermuda. Commonwealth Agricultural Bureaux, 1959

[Conyers, J. & Wingate, D. 1996. *Eumeces longirostris*. In: IUCN 2010. IUCN Red List of Threatened Species.](#)

Summary: Available from: <http://www.iucnredlist.org/apps/redlist/details/8218/0> [Accessed 22 June 2010]

Davenport, John; Jeremy Hills; Anne Glasspool and Jack Ward, 2001. Threats to the Critically Endangered endemic Bermudian skink *Eumeces longirostris*. *Oryx* Vol 35 No 4 October 2001

Gorman, George C., Y. J. Kim, S. Y. Yang, 1978. The Genetics of Colonization: Loss of Variability among Introduced Populations of *Anolis* Lizards (Reptilia, Lacertilia, Iguanidae). *Journal of Herpetology*, Vol. 12, No. 1 (Feb. 27, 1978), pp. 47-51

Losos, Jonathan B., 1996. Dynamics of Range Expansion by Three Introduced Species of *Anolis* Lizards on Bermuda. *Journal of Herpetology*, Vol. 30, No. 2 (Jun., 1996), pp. 204-210

Macedonia, Joseph M. and David L. Clark, 2003. Headbob Display Structure in the Naturalized *Anolis* Lizards of Bermuda: Sex, Context, and Population Effects. *Journal of Herpetology*, Vol. 37, No. 2, pp. 266-276, 2003

Nicholson, Kirsten E.; Glor, Richard E.; Kolbe, Jason J.; Larson, Allan; Hedges, S. Blair; Losos, Jonathan B., 2005. Mainland colonization by island lizards. *Journal of Biogeography*. 32(6). JUN 05. 929-938

[Reptiles Database, 2010. *Anolis grahami* Gray, 1845](#)

Summary: Available from: <http://reptile-database.reptarium.cz/species.php?genus=Anolis&species=grahami> [Accessed September 8 2010]

Schoener, Thomas W., 1975. Presence and Absence of Habitat Shift in Some Widespread Lizard Species. *Ecological Monographs*, Vol. 45, No. 3 (Summer, 1975), pp. 233-258

Strong, D., B. Leatherman, and B.H. Brattstrom. 1993. Two new methods for catching small fast lizards. *Herpetological Review* 24:22-23.

Wingate, David B., 1965. Terrestrial Herpetofauna of Bermuda. *Herpetologica*, Vol. 21, No. 3 (Sep. 24, 1965), pp. 202-218