

# **GLOBAL INVASIVE SPECIES DATABASE**

FULL ACCOUNT FOR: Berberis darwinii

Berberis darwinii System: Terrestrial

Kingdom	Phylum	Class	Order	Family
Plantae	Magnoliophyta	Magnoliopsida	Ranunculales	Berberidaceae

**Common name** Darwin's Berberis (English), Darwin's Barberry (English), berbére (French)

**Synonym** 

**Similar species** Berberis thunbergii, Berberis vulgaris

**Summary** Berberis darwinii exists as varying habits in New Zealand from shrubs with

interlaced branches, reaching 3-4m in height and 3-6m wide in the open and at the forest edge to lianoid small trees growing to about 10m and spreading 15m under the forest canopy. Dispersal is key to *B. darwinii*'s survival. Introduced bird species act as dispersers in New Zealand. Despite being considered a shade-tolerant species it has been shown that *B. darwinii* actually requires high light environments to germinate. It is tolerant of both

frost and drought.



view this species on IUCN Red List

### **Species Description**

Berberis darwinii exists as varying habits in New Zealand, from shrubs with interlaced branches, reaching 3-4m in height and 3-6m wide in the open and at the forest edge to lianoid small trees growing to about 10m and spreading 15m under the forest canopy (Allen & Wilson, 1992). Dispersal is key to Berberis darwinii's survival as shown by MacAlpine & Jesson (2008). They found great numbers of seeds dispersed at least 150m, with others also being consistantly detected up to 450m away from the parent population. This was further shown in their study with seedling survival; nearly all seedlings under the parent population died within the first 5 months of germination. Despite being considered a shade-tolerant species (Webb et al 1988), MacAlpine & Jesson (2008) and MacAlpine et al (2008) have shown that B. darwinii actually requires high light environments to germinate. It is tolerant of both frost and drought (Allen 1991; Timmins & Mackenzie 1995; as seen in McAlpine & Jesson, 2007)

### **Lifecycle Stages**

Berberis darwinii flowers in Spring (August to November) and fruits in Summer (December to February) in New Zealand, (MacAlpine & Jesson, 2008), although fruits and flowers can both be seen on the species nearly year-round (Webb *et al*, 1988; as seen in McAlpine & Jesson, 2008).

#### Reproduction

The main pollinator in New Zealand of Berberis darwinii is the honey bee (Apis mellifera) (Allen & Wilson, 1992).



# **GLOBAL INVASIVE SPECIES DATABASE**

FULL ACCOUNT FOR: Berberis darwinii

### **Management Info**

<u>Preventative measures</u>: On the Falkland Islands, Whitehead (2008; as seen in Oyley *et al*, 2008) conducted a risk assessment on potentially invasive species. *Berberis darwinii* scored 18. Any species with a score over 15 is considered invasive.

Physical: Several management techniques have been trialled on *B. darwinii* in New Zealand (Ward & Henzell, 2004), each a combination of physical and chemical methods. Cut-Stem treatment on saplings proved very effective and involves the cutting of all sapling stems coupled with an application of herbicide. A Wedge-Method and a Trunk Injection method, whilst not entirely physical, each proved very effective when also coupled with herbicide. The Wedge-Method involves the cutting of wedges out of the trees at constant intervals and applying herbicide within each wedge. The trunk Injection technique involves the drilling of holes into the main tree trunks and injecting herbicide within each hole. The Cut-Stem and Wedge-Method each had a 100% kill rate in trials, whereas the Trunk Injection had an 80-90% kill rate. These methods have only been trialled within New Zealand (Ward & Henzell, 2004).

Chemical: Several management techniques have been trialled on *B. darwinii* in New Zealand (Ward & Henzell, 2004), each a combination of physical and chemical methods. Cut-Stem treatment on saplings proved very effective and involves the cutting of all sapling stems coupled with an application of herbicide. A Wedge-Method and a Trunk Injection method, each proved very effective when also coupled with herbicide. The Wedge-Method involves the cutting of wedges out of the trees at constant intervals and applying herbicide within each wedge. The trunk Injection technique involves the drilling of holes into the main tree trunks and injecting herbicide within each hole. The Cut-Stem and Wedge-Method each had a 100% kill rate in trials, whereas the Trunk Injection had an 80-90% kill rate. The most effective herbicide found within these trials was the Vigilant herbicide (a mixture of 5% picloram potassium salt). These methods have only been trialled within New Zealand (Ward & Henzell, 2004).

Biological: MacAlpine et al (2008) suggest that controlling efforts should be directed at removing individuals in high light as opposed to across all environments, due to its fast growth capabilities in such an environment. MacAlpine & Jesson (2008) further suggest that control should focus on elimination of the seed source, however when this is not possible, efforts should ignore seedlings under parent populations as they generally don't survive anyway, and that the seed bank should be of no concern as most seeds do not last for more than 1 year.

### **Principal source:**

**Compiler:** IUCN SSC Invasive Species Specialist Group (ISSG) with support from the Overseas Territories Environmental Programme (OTEP) project XOT603, a joint project with the Cayman Islands Government - Department of Environment

Review:

**Pubblication date: 2010-06-08** 

**ALIEN RANGE** 

[1] AUSTRALIA [1] FALKLAND ISLANDS (MALVINAS)

[1] IRELAND [1] NEW ZEALAND

[3] UNITED STATES

**BIBLIOGRAPHY** 

16 references found for Berberis darwinii

**Managment information** 



# **GLOBAL INVASIVE SPECIES DATABASE**

FULL ACCOUNT FOR: Berberis darwinii

IUCN/SSC Invasive Species Specialist Group (ISSG)., 2010. A Compilation of Information Sources for Conservation Managers.

**Summary:** This compilation of information sources can be sorted on keywords for example: Baits & Lures, Non Target Species, Eradication, Monitoring, Risk Assessment, Weeds, Herbicides etc. This compilation is at present in Excel format, this will be web-enabled as a searchable database shortly. This version of the database has been developed by the IUCN SSC ISSG as part of an Overseas Territories Environmental Programme funded project XOT603 in partnership with the Cayman Islands Government - Department of Environment. The compilation is a work under progress, the ISSG will manage, maintain and enhance the database with current and newly published information, reports, journal articles etc.

Otley H, Munro G, Clausen A and Ingham B. 2008. Falkland Islands State of the Environment Report 2008. Falkland Islands Government and Falklands Conservation, Stanley.

Ward, B. G. and R.F. Henzell, 2001. Use of herbicidal gels on woody weeds. DOC Science Internal Series 162

Summary: Available from: http://www.doc.govt.nz/upload/documents/science-and-technical/dsis162.pdf [Accessed July 3 2010]

#### **General information**

Alien Plants in Ireland, 2010. Berberis darwinii Hook.

**Summary:** Available from: http://www.biochange.ie/alienplants/result\_options.php?species=51&families=Berberidaceae&p=i&blz=1 [Accessed July 3 2010]

Allen, R. B., 1991. A preliminary assessment of the establishment and persistence of *Berberis darwinii* Hook., a naturalised shrub in secondary vegetation near Dunedin, New Zealand. New Zealand Journal of Botany, 1991, Vol. 29: 353-360

Allen, R. B. and W. G. Lee, 2001. Woody weed dispersal by birds, wind and explosive dehiscence in New Zealand. New Zealand Plant Protection 54:61-66 (2001)

Summary: Available from: http://www.nzpps.org/journal/54/nzpp\_540610.pdf [Accessed July 3 2010]

Allen, R. B. & J. B. Wilson, 1992. Fruit and seed production in *Berberis darwinii* Hook., a shrub recently naturalised in New Zealand. New Zealand Journal of Botany, 1992, Vol. 30: 45-55

Dallimore, W., 1919. The Falkland Islands. Forestry. Tussock Grass. Bulletin of Miscellaneous Information (Royal Gardens, Kew), Vol. 1919, No. 5 (1919)

Integrated Taxonomic Information System (ITIS), 2010. Berberis darwinii Hook.

**Summary:** Available from: http://www.itis.gov/servlet/SingleRpt/SingleRpt?search\_topic=TSN&search\_value=506843 [Accessed July 3 2010] McAlpine, Kate G. and Linley K. Jesson, 2007. Biomass allocation, shade tolerance and seedling survival of the invasive species *Berberis darwinii* (Darwin s barberry). New Zealand Journal of Ecology (2007) 31(1): 1-12

McAlpine, Kate G. and Linley K. Jesson, 2008. Linking seed dispersal, germination and seedling recruitment in the invasive species *Berberis darwinii* (Darwin s barberry). Plant Ecol (2008) 197:119 129

McAlpine, Kate G., Linley K. Jesson and David, S. Kubien, 2008. Photosynthesis and water-use efficiency: A comparison between invasive (exotic) and non-invasive (native) species. Austral Ecology (2008) 33, 10 19

Please refer this to the updated Varnham, there is a secondary ref to the Guernsey Botanical Records

USDA, ARS, 2010. Taxon: Berberis darwinii Hook. USDA, ARS, National Genetic Resources Program. Germplasm Resources Information Network - (GRIN) [Online Database].

Summary: Available from: http://www.ars-grin.gov/cgi-bin/npgs/html/taxon.pl?6845 [Accessed July 3 2010]

USDA, NRCS, 2010. Berberis darwinii Hook. Darwin s berberis. The PLANTS Database. National Plant Data Center, Baton Rouge, LA 70874-4490 USA.

**Summary:** Available from: http://plants.usda.gov/java/profile?symbol=BEDA [Accessed July 3 2010]

Williams, Peter A., 2006. The role of blackbirds (*Turdus merula*) in weed invasion in New Zealand. New Zealand Journal of Ecology (2006) 30(2): 285-291

Summary: Available from: http://www.newzealandecology.org.nz/nzje/free\_issues/NZJEcol30\_2\_285.pdf [Accessed July 3 2010]