

Cynara cardunculus

System: Terrestrial

Kingdom	Phylum	Class	Order	Family
Plantae	Magnoliophyta	Magnoliopsida	Asterales	Asteraceae
Common name	Artischocke (German), artichaut commun (French), Scottish thistle (English), wild cardoon (English), cynara (English), Scotch thistle (English), globe artichoke (English), artichoke thistle (English), Spanish thistle artichoke (English), wild artichoke (English), cardoon (English), cardon d'Espagne (French), Kardone (German), alcachofa (Spanish), alcachofra (Portuguese), alcaucil (Spanish), cardo de comer (Spanish), cardo (Portuguese), cardo (Spanish), Gemüseartischocke (German)			
Synonym	Cynara cardunculus , L. var. altilis Cynara cardunculus , L. var. eu-cardunculus Hayek Cynara cardunculus , L. var. ferocissima Lowe Cynara cardunculus , L. var. hortensis Brotero Cynara cardunculus , L. var. inermis Cynara cardunculus , L. var. sativaBrotero Cynara cardunculus , L. var. sativa Moris Cynara cardunculus , L. var. sativa Moris Cynara cardunculus , L. var. soclymus Cynara cardunculus , L. var. soclymus Cynara cardunculus , L. var. soclymus Cynara cardunculus , L. typica Willkomm ex. Mariz Cynara corsica , Viviani Cynara esculenta Cynara ferox , Tenore fide Steudel Cynara hortesis Cynara scolymus , L. var. hortensis Cynara scolymus , L. var. hortensis Cynara scolymus , L. var. minor Brotero Cynara scolymus , L. var. minor Brotero Cynara scolymus , L. var. mutica de Visiani Cynara scolymus , L. var. mutica de Visiani Cynara sylvestris , de Lamarck Cynara sylvestris , Lam. var. spinosus de Lamarck			
Similar species				
Summary	Artichoke thistle (<i>Cynara cardunculus</i>) is a large perennial herb native to the Mediterranean region of Europe. It grows rapidly and produces very deep roots, and forms large rosettes up to 1 m or more in diameter. It has invaded Australia, California and parts of South America where it is an aggressive weed, capable of forming dense, massive monospecific stands to the exclusion of all other plant species.			
C REP	view this sp	pecies on IUCN Red List		



FULL ACCOUNT FOR: Cynara cardunculus

Species Description

Cynara cardunculus is a perennial herb which reaches 1.1 to 1.8 m in height (Wiklund, 1992). It is often richly branched and produces a very deep perennial taproot from which the plant regenerates each year (Kelly & Pepper, 1996). The root crown can also produce multiple plants, serving as a form of asexual reproduction. *C. cardunculus* forms a rosette up to 1 m or more in diameter composed of very large leaves. Leaves are silky greyish-green on the upper surface and appear almost white on the under surface because of the dense mat of white hairs. Leaves are deeply divided, each lobe ending in a strong yellow spine (DAF, n.d.). In spring a tall, branching, flowering stem up to 2 m high is produced. One large, fragrant blue or purple flower forms at each branch apex. The flower heads consist of tubular florets surrounded by a series of rigid bracts (phyllaries) tipped with spines. Each plant may produce up to 50 heads. Seeds are brown to black, about 5 mm long with a smooth covering. Seeds are wind-dispersed, each with a "parachute" of feathery bristles (pappus) about 4 cm long (DAF, n.d.). Wild representatives of *C. cardunculus* may be distinguished from other *Cyanara* species by their deeply dissected leaves and many long spines (Wiklund, 1992).

Notes

Cynara cardunculus has been divided into two subspecies: the western morph is subsp. *flavescens*; the eastern morph is subsp. *cardunculus* (Wiklund, 1992). The subspecies differ mainly by a yellowish margin on the middle involucral bract (Wiklund, 1992). There are also a number of varieties of *C. cardunculus* including the cultivated artichoke var. *scolymus* (L.) Fiori, the wild artichoke var. *sylvestris* (Lam.) Fiori and the leafy cardoon var. *altilis* DC (Pignone & Sonnante, 2004).

Lifecycle Stages

Cynara cardunculus is a perennial herb that may have a lifecycle exceeding 10 years (Archontoulis *et al.*, 2010). Seeds are set in summer, germinate in autumn and rapidly form a large flat rosette of leaves and a deep fibrous tap root (DAF, n.d.). Two phases of growth are distinguished which comprise a) vegetative stage which has a duration of approximately 220 days (from September to April) and b) the reproductive phase (duration 95 days; from May to August) (Archontoulis *et al.*, 2010). In the northern hemisphere flowering occurs mainly in June and July, although flowering has been observed between May and August. In the southern hemisphere flowering has been observed between November to January (Wiklund, 1992).

The first set of rosette leaves normally dies off over the summer and new leaves are formed in the following autumn, which resprout from the perennial tap root (DAF, n.d.).

Uses

Cynara cardunculus can be used as forge in winter and spring, as a raw material in the paper pulp industry, and most importantly as a solid and liquid biofuel in the bio-energy sector in its native Mediterranean range (Curt *et al.*, 2004). It requires little water, nitrogen fertilisers and agrochemicals due to its rapid growth and deep root system. It also improves soil characteristics and prevents erosion. Considering the modest inputs needed to grown *C. cardunculus* it may produce the cheapest biofuel compared to all other bio-energy crops known (Grammelis *et al.*, 2008).

The bases of the phyllaries and the fleshy receptacle of *C. cardunculus* are edible. The petioles and roots may also be edible if properly prepared (Kelly & Pepper, 1996).

The artichoke has also been used for medicinal purposes (Adly, 1985 in Wiklund, 1992). It is a source of inulin, cynarin and silymarin (Curt *et al.*, 2002 in Grammelis *et al.*, 2008). The latter two are bitter-tasting compounds, which are found in the leaves, and improve liver and gall bladder function, and stimulate the secretion of digestive juices, especially bile, and lowers blood cholesterol levels (Grammelis *et al.*, 2008). Dried florets of artichoke thistle have been used to coagulate milk (Mathews, 1830 in Wiklund, 1992).



FULL ACCOUNT FOR: Cynara cardunculus

Habitat Description

In California *Cynara cardunculus* is a particularly problematic pest plant on poorly managed and overgrazed rangelands. It can also invade relatively undisturbed natural habitats such as coastal sage scrub, chaparral and riparian woodlands (Kelly & Pepper, 1996).

The species can tolerate harsh environments with high temperatures and water stress, even on thin and unproductive soils, due to its deep root system which may exceed 5 m in depth (Archontoulis *et al.*, 2010). It appears to favour soils with a high clay content, but can also tolerate coarse-textured soils (Thomsen *et al.*, 2004).

Reproduction

Although *C. cardunculus* reproduces vegetatively from the roots and crown of mature plants, the species is thought to spread primarily by seed (Marushia & Holt, 2006). The plant occasionally flowers in the first year, but usually not until the second summer (DAF, n.d.). Each plant produces from 1 to 50 inflorescences per rosette per year (Marushia & Holt, 2006). Flowers have strong fragrance and produce abundant nectar that attracts a wide variety of insects which carry out pollination (Foury, 1967 in Wiklund, 1992). Up to 600 achenes (seeds) are produced per inflorescence, which are equipped with a large pappus (Marushia & Holt, 2006) for dispersal via wind (Bowler, 2008). However, in contrast to most wind-dispersed plants *C. cardunculus* has larger seeds (20-60 mg) and the pappus often becomes detached before dispersal. (Parsons, 1973, Hillyard, 1985 in Marushia & Holt, 2006). Seeds travel further in non-vegetated sites (more than 40 m) compared to vegetated sites (less than 20m) where seeds become trapped by other plants (Marushia & Holt, 2006). However most seeds disperse within 3 m of source plants (Marushia & Holt, 2006).\r\n

Flowers and rosettes senesce in late summer; the perennial tap root persists and plants regenerate from this the following spring (Wiklund, 1996; Kelly & Pepper, 1996)

General Impacts

<u>Competition</u>: The artichoke thistle forms large monocultures where it constitutes 100% of the vegetation cover, completely excluding most other plant species. It has been known to reach densities of over 20,000 plants per acre (Kelly & Pepper, 1996). They produce large rosettes (up to 2 m diameter) and produce an expansive litter layer each year that exclude other plants via shading or mechanical impedance, or by production of allelochemicals. Seedlings of *C. cardunculus* have high tolerance for intraspecific competition, and will grow at high densities at the margins of adult rosettes. This is likely to be a factor allowing high density monocultures (Marushia & Holt, 2008). These large monocultures result in fragmentation of higher quality habitat (Kelly & Pepper, 1996). Native forbs are particularly vulnerable to competition with *C. cardunculus* (White & Holt, 2005).\r\n\r\n

<u>Reduction of native biodiversity</u>: Invasion of the fragile canyon ecosystem in California has led to reduction of populations of the endangered San Diego thorn mint (*Acanthimentha ilicifolia*). The absence of wildlife trails in artichoke monocultures indicates that infestations are an obstacle to wildlife movement (Kelly & Pepper, 1996). However Bowler (2008) report that some animals can utilise artichoke: squirrels and birds forage on seeds, it provides high perch sites for birds, and some hummingbird species have high nesting success on artichoke branches and leaves.\r\n

<u>Agricultural</u>: Artichoke thistle stands can become so thick that forage production and livestock movement are impeded (Thomsen *et al.*, 2004).

<u>Modification of nutrient regime</u>: *C. cardunculus* produces greater aboveground biomass than native plants, which correspondingly increases midday ecosystem carbon assimilation and evapotranspiration rates. Additionally, litter quality and nitrogen content are increased in areas with *C. cardunculus*. The flow-on effects of such changes are unresolved but may affect carbon cycling, soil organic carbon quantities and increase soil respiration rates (Potts *et al.*, 2008). *C. cardunculus* may also affect soil microbial communities and nutrient and water feedback cycles, although this has not yet been investigated (Potts *et al.*, 2008).\r\n <u>Human health</u>:The plant is reported to cause contact dermatitis in some people (Victoria Resources Online, 2009).



FULL ACCOUNT FOR: Cynara cardunculus

Management Info

<u>Physical control</u>: The artichoke thistle is large and has a perennial tap-root capable of regeneration, unless the entire root system has been destroyed. Thus attempts to control this plant by plowing, chaining, scraping and bulldozing are often unsuccessful (Kelly & Pepper, 1996). Mowing is unsuccessful as the shoot apical meristem is buried below the soil surface (White & Holt, 2005). However, intensive manual pulling by large numbers of people have been successful in parts of California (Thomsen *et al.*, 2004).\r\n

<u>Chemical control</u>: Foliar application of 2% glyphosate (Round-Up, Monsanto) applied during the active growth period (January-July) was successful in controlling artichoke thistle in Los Peñasquitos Canyon Preserve, California. The herbicide should cover the plant's leaf surface to ensure adequate application. Young seedlings can be killed with one application. Most seedlings occur at or near the margins of adult plants; therefore, spraying a 1 m buffer zone around adults will increase the long-term effectiveness of an herbicide spray treatment by removing the majority of seedlings (Marushia & Holt, 2008). In larger plants the greatest success rate was achieved by applying herbicide while the flower stalk was bolting (Kelly & Pepper, 1996). In areas where spraying is not advisable due to sensitive habitat, herbicide can be brushed on uncut rosette leaves (Tallis & White, 2009).\r\n

<u>Seeding</u>:Once artichoke has been removed from a site, revegetation with natives is relatively straightforward, as artichoke stands rarely contain a large seedbank. Seeding with native plant species has been successful at a number of sites in California. However if glyphosphate was used a herbicide to eradicate artichoke, seeds should not be introduced for at least 1.5 months (Bowler, 2008). Because root competition strongly affects artichoke thistle, the ideal species for revegetation are those with rapidly developing root systems, and should include a combination of shallow and deep rooted species to maximise competition (White & Holt, 2005).\r\n Biological control: Biocontrol is not an option due to the close phylogenetic relationship to the cultivated artichoke (*C. skolymus*) (Kelly & Pepper, 1996; Thomsen *et al.*, 2004).

Principal source:

Compiler: National Biological Information Infrastructure (NBII) & IUCN/SSC Invasive Species Specialist Group (ISSG)

Review: Dr. Robin Marushia, Associate Director, Koffler Scientific Reserve, Ontario

Pubblication date: 2011-02-23

ALIEN RANGE

[2] AUSTRALIA[1] CHILE[1] NEW ZEALAND[1] SOUTH AMERICA[1] UNITED KINGDOM

[1] CENTRAL AMERICA
[1] ECUADOR
[1] NORTHERN EUROPE
[1] SPAIN
[4] UNITED STATES

BIBLIOGRAPHY

24 references found for Cynara cardunculus

Managment information

Bowler, Peter A., 2008. Artichoke as an Ecological Resource and Its Utility as a Precursor to Restoration (California). March 2008 Ecological Restoration 26:1

Summary: Available from: http://er.uwpress.org/cgi/reprint/26/1/7.pdf [Accessed 17 July 2010]

Crosti, Roberto; Carmela Cascone and Salvatore Cipollaro, 2010. Use of a weed risk assessment for the Mediterranean region of Central Italy to prevent loss of functionality and biodiversity in agro-ecosystems. Biol Invasions (2010) 12:1607 @1616

Department of Agriculture and Food, Governement of Western Australia, Declared plant in Western Australia Artichoke thistle (Cynara cardunculus)

Summary: Available from: http://www.agric.wa.gov.au/objtwr/imported_assets/content/pw/weed/decp/artichoke_thistle.pdf [Accessed 17 July 2010]



FULL ACCOUNT FOR: Cynara cardunculus

Department of Primary Industries, 2009. Artichoke thistle Impact Assessment (*Cynara cardunculus*) in Victoria. Victorian Resources Online. Government of Victoria

Summary: Available from:

http://www.dpi.vic.gov.au/dpi/vro/vrosite.nsf/0d08cd6930912d1e4a2567d2002579cb/6e3a7a1eae8154e7ca25760800822394/\$FILE/Artichok e%20thistle%20Impact%20Assessment%20(Cynara%20cardunculus).pdf [Accessed 17 July 2010]

Kelly, Mike and Alan Pepper, 1996. Controlling Cynara cardunculus (Artichoke Thistle, Cardoon, etc.). California Exotic Pest Plant Council 1996 Symposium Proceedings

Summary: Available from: http://www.cal-ipc.org/symposia/archive/pdf/1996_symposium_proceedings1817.pdf [Accessed 17 July 2010] Potts, Daniel L.; Harpole, W. Stanley; Goulden, Michael L.; Suding, Katharine Nash, 2008. The impact of invasion and subsequent removal of an exotic thistle, *Cynara cardunculus*, on CO2 and H2O vapor exchange in a coastal California grassland. Biological Invasions. 10(7). OCT 2008. 1073-1084.

Tallis, J & White, V., 2009. Using Herbicide to control artichoke thistle in rare plant habitat. CNPS 2009 Conservation Conference: Strategies and Solutions. [np]. 2009

Summary: Control of invasive non-native plants is a challenge to land managers worldwide. This challenge is magnified when non-natives plants invade rare plant habitat because the tools and techniques used to eradicate one species may negatively impact an adjacent protected species. As a means of developing a method of controlling artichoke thistle (Cynara cardunculus) growing on serpentine soils near San Jose, California, this experiment tested two different herbicides and two different application approaches. Our treatments included applying Roundup Pro registered or Transline registered by brushing herbicide on cut stems or uncut leaves. Our results showed that both herbicides and both application techniques were significantly effective at killing artichoke thistle during the 7-month experiment running from 2007-2008. This result suggests that land managers can use a less expensive herbicide (Roundup Pro registered) and apply it with a less time consuming method (brushing on uncut leaves) when controlling artichoke thistle in conditions unsuitable for spraying

General information

Archontoulis, S. V.; Struik, P. C.; Vos, J.; Danalatos, N. G., 2010. Phenological growth stages of *Cynara cardunculus*: codification and description according to the BBCH scale. Annals of Applied Biology. 156(2). 2010. 253-270.

Charles Darwin Foundation (CDF), 2010. Cynara cardunculus: The Charles Darwin Foundation Galapagos Species Checklist **Summary:** Available from: http://www.darwinfoundation.org/datazone/checklists/vascular-plants/magnoliophyta/cynara-cardunculus-l/ [Accessed 17 July 2010] Curt, M. D., G. S@nchez and J. Fern@ndez, 2002. The potential of Cynara cardunculus L. for seed oil production in a perennial cultivation

Curt, M. D., G. Sonchez and J. Fernondez, 2002. The potential of *Cynara cardunculus* L. for seed oil production in a perennial cultivation system. Biomass and Bioenergy Volume 23, Issue 1, July 2002, Pages 33-46

Grammelis, Panagiotis; Anastasia Malliopoulou; Panagiotis Basinas and Nicholas G. Danalatos, 2008. Cultivation and Characterization of Cynara cardunculus for Solid Biofuels Production in the Mediterranean Region. Int. J. Mol. Sci. 2008, 9, 1241-1258; DOI: 10.3390/ijms9071241

Summary: Available from: http://www.mdpi.org/ijms/papers/i9071241.pdf [Accessed 17 July 2010]

Integrated Taxonomic Information System (ITIS), 2010. Cynara cardunculus L.

Summary: Available from: http://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=37221 [Accessed 17 July 2010] Marushia, R. and J. S. Holt. 2003. Patterns of seedling establishment in artichoke thistle, *Cynara cardunculus*. Proc. Calif. Invasive Plant Council, Annual Symposium, Vol. 9, p. 59-60.

Council, Annual Symposium, Vol. 9, p. 59-60. Marushia, R. and J. S. Holt. 2004. Seed dispersal and patterns of seedling establishment in *Cynara cardunculus*. Ecol. Soc. Amer. Abstr., Ann. Meeting, Available on CD.

Marushia, Robin G.; Holt, Jodie S., 2006. The effects of habitat on dispersal patterns of an invasive thistle, *Cynara cardunculus*. Biological Invasions. 8(4). JUN 2006. 577-593.

Marushia, Robin G.; Holt, Jodie S., 2008. Reproductive strategy of an invasive thistle: effects of adults on seedling survival. Biological Invasions. 10(6). AUG 2008. 913-924.

Summary: Available from:

http://faculty.rio.edu/dalthoff/Courses/ISBEM/Set%201/TP-Marushia%20and%20Holt-Invasive%20Thistle-2008-BI.pdf [Accessed 17 July 2010] Perry, Roger, 1984. Juan Fern@ndez Islands: A Unique Botanical Heritage. Environmental Conservation, 11, pp 72-76

Summary: Available from: http://www.montsevila.org/PapersSCI/Vila_etalJVS.pdf [Accessed 17 July 2010] Pignone, Domenico and Gabriella Sonnante, 2004. Wild artichokes of south Italy: did the story begin here? Genetic Resources and Crop Evolution 51: 577\$580, 2004.

Summary: Available from: http://www.montsevila.org/PapersSCI/Vila_etalJVS.pdf [Accessed 17 July 2010]

Thomsen, Craig D., G. Douglas Barbe, William A. Williams & Melvin R. George, 1986. Escaped artichokes are troublesome pests. California Agriculture, March-April 1986

Summary: Available from: http://ucce.ucdavis.edu/files/repositoryfiles/ca4003p7-62935.pdf [Accessed 17 July 2010]

USDA, ARS, 2010. Taxon: Cynara cardunculus L. National Genetic Resources Program. Germplasm Resources Information Network - (GRIN) [Online Database]. National Germplasm Resources Laboratory, Beltsville, Maryland.

Summary: Available from: http://www.ars-grin.gov/cgi-bin/npgs/html/taxon.pl?12839 [Accessed 17 July 2010] USDA-NRCS, 2010. Cynara cardunculus L. cardoon. The PLANTS Database (http://plants.usda.gov, 4 August 2010). National Plant Data

Center, Baton Rouge, LA 70874-4490 USA. Summary: Available from: http://plants.usda.gov/java/profile?symbol=CYCA [Accessed 17 July 2010]

Vila, Montserrat; Pino, Joan; Montero, Ana; Font, Xavier, 2010. Are island plant communities more invaded than their mainland counterparts? Journal of Vegetation Science. 21(3). JUN 2010. 438-446.

Summary: Available from: http://www.montsevila.org/PapersSCI/Vila_etalJVS.pdf [Accessed 17 July 2010]

White, Virginia A.; Holt, Jodie S., 2005. Competition of artichoke thistle (*Cynara cardunculus*) with native and exotic grassland species. Weed Science. 53(6). NOV-DEC 2005. 826-833.

Wiklund, Annettte, F.L.S, 1992. The genus Cynara L. (Asteraceae-Cardueae). Botanical Journal of the Linnean Society (1992), 109: 75-123.