

**GLOBAL INVASIVE SPECIES DATABASE** 

FULL ACCOUNT FOR: Duttaphrynus melanostictus

#### Duttaphrynus melanostictus

System: Freshwater\_terrestrial

Kingdom	Phylum	Class	Order	Family	
Animalia	Chordata	Amphibia	Anura	Bufonidae	
Common name	Asian common toad (English)				
Synonym	Bufo melanostictus ,Schneider, 1799				
Similar species					
Summary	The species Duttaphrynus melanostictus is one of the most widespread species in its genus across tropical Southeast Asia. The species prays on native insects and other small vertebrate species. It causes decile in population size of reptiles and amphibians through completion and is a risk to human health due to its toxicity. The species is listed in the IUCN Red List of Threatened Species as Least Concern (IUCN, 2018).				
Synonym Similar species Summary	Bufo melan The species species in in native inser population human hea Threatened	Bufo melanostictus ,Schneider, 1799 The species Duttaphrynus melanostictus is one of the most widesprea species in its genus across tropical Southeast Asia. The species prays native insects and other small vertebrate species. It causes decile in population size of reptiles and amphibians through completion and is a human health due to its toxicity. The species is listed in the IUCN Red Threatened Species as Least Concern (IUCN, 2018).			



view this species on IUCN Red List

#### **Species Description**

The colour of D. melanostictus is greyish or reddish brown. The size adults range from 60 to 99mm The body size of females is slightly larger than the body size of males (Berry & Bullock, 1962). The head has elevated bony ridges, with long dark crests that border the eyelids and run down on either side of the eye. Tadpoles are black and small, up to 15 mm long (Csurhes, 2016). The species is widely spread in Southeast Asia. It commonly occurs in northern Pakistan through Bangladesh, India, Sri Lanka, southern China, Myanmar, Laos, Viet Nam, Thailand, Cambodia to Malaysia, Singapore and parts of Indonesia.

#### Notes

According to species distribution models the species is potentially invasive to Madagascar and Australia (Csurhes, 2016; Moore et al., 2015). Individuals have never been recorded on the Australian mainland. However few individuals have been detected on shipping containers by Australian Quarantine and Inspection Service (AQIS) (Trainor, 2009).

#### **Lifecycle Stages**

The larvae metamorphosis happens within 25 to 30 days. The length of the larvae varies from 34 to 90 days. The average life span is 4 to 10 years (Berry & Bullock, 1962).

#### Uses

The species is introduced unintentional and intentional. In Timor-Leste the species was introduced to control mosquito numbers (Trainor, 2009).

#### **Habitat Description**

The species occurs naturally in open habitats such as grasslands and deciduous dipterocarp savannah but also in the proximity of human habitation and cultivation (Berry & Bullock, 1962; Wogan et al., 2016). It occurs up to a sea level of 2000m in temperate, tropical and sub-tropical habitats. The species can tolerate a salinity level of up to 1% (McClelland et al, 2015).



**GLOBAL INVASIVE SPECIES DATABASE** 

FULL ACCOUNT FOR: Duttaphrynus melanostictus

#### Reproduction

The species is oviparous. The Asian common toad (D. melanostictus) breeds twice per year, most likely once per year, depending on rainfall and temperature. The breeding kates place in still water bodies or slowly moving river streams. The breeding season in semitropical regions in Asia dues from February to April (Huang et al., 1997). "Males congregate at breeding sites where they call to females. The call is a 'creo-o-o'; cro-ro-ro-ro-ro-ro' that is repeated in choruses rather monotonously. The calling males may be heard throughout the night during rain and sometimes on overcast days", (Daniels 2005: in Csurhes, 2016). Approximately 40,000 eggs are laid per clutch.

#### Nutrition

D. melanostictus mainly feeds on insects within a size rage of 5.0 to 20.0 mm (Berry & Bullock, 1962).

#### **General Impacts**

The species has been introduced to Maluka, Papua and Sulawesi, reported in Timor Leste and Madagascar. The Asian common toad has been intercepted at Australian borders. Impact on human health: The species is toxic. Contact with the toxins in parotid glands on its back can cause allergic reactions, because of several bioactive compounds with lethal, hypotensive, hypertensive, neurotoxic, cardio toxic, hemolytic and sleep inducing factors (Das et al. 2000). In Timor-Leste the species became a public health issue, because villagers, who's diet include native toads, have poisoned themselves eating the toxic alien toad species. The consumption of the species can cause serious illness and sometimes death (McCelland et al, 2015). In Laos the death of a 6 year old boy was reported after he had consumed a dish containing grilled Asian common toad. Several other cases were reported, where patients suffered from severe vomiting few hours after eating the species. Toad poisonning is serious impact on human heath in the rural area of Laos (Keomany et al., 2007). Impact on biodiversity: In Madagascar the Asian common toad is currently spreading rapidly. There are fears that it could have severe ecological impacts on amphibian prey species (van Dijk et al., 2004). In areas where the species established the species is impacting native species richness. Several native species populations declined after the arrival of D. melanostictus (Trainor, 2009). Small native species are impacted by direct toad predation. The species prays on populations of small isect groups, scincid lizards, geckos and land snails (Trainor, 2009). The impact of D. melanostictus on livestock is unknown.

#### **Management Info**

The spread of the species is influenced by climate, availability of wetland sites for breeding, daytime shelter and food. The population's increse is positivley correlated to the rainfall (Trainor, 2009). Early detection mesures: It is recommanded to implement quarantine procedures in all regions where the species could potentially invade(Trainor, 2009). Andreone (2014) recommands to use genetic analyses to identify the source of introduction and will screen toads for pathogens and parasites. Public education and awareness: It is of high impotance to increse the public awareness to ensure the identification of the species. Eradication actions need to be carefully considered in order to protect native tead populations in the same place (Mecke, 2014). Larvae of D. melanostictus, for example, survive in streams, puddles and brackish waters (Mecke, 2014).

Principal source: Csurhes, 2016. Berry & Bullock, 1962. Trainor, 2009.

**Compiler:** 

**Review:** 

#### **Pubblication date:**

#### ALIEN RANGE

[1] AUSTRALIA[1] BRUNEI DARUSSALAM[2] INDONESIA[1] MADAGASCARGlobal Invasive Species Database (GISD) 2025. Species profile Duttaphrynus melanostictus.Available from: <a href="https://www.iucngisd.org/gisd/species.php?sc=1904">https://www.iucngisd.org/gisd/species.php?sc=1904</a> [Accessed 05 July 2025]



# **GLOBAL INVASIVE SPECIES DATABASE**

FULL ACCOUNT FOR: Duttaphrynus melanostictus

# [1] PAPUA NEW GUINEA[1] UNITED ARAB EMIRATES

[1] TIMOR-LESTE [1] UNITED STATES

### Red List assessed species 3: LC = 3;

Ingerophrynus biporcatus LC Varanus salvator LC Trimeresurus insularis LC

## 22 references found for Duttaphrynus melanostictus

#### Managment information

**BIBLIOGRAPHY** 

wa.gov.au (2017). https://www.agric.wa.gov.au/amphibians-and-reptiles/biosecurity-alert-asian-black-spined-toad?nopaging=1 Andreone, F. (2014). Madagascar: Risk review is under way for invasive toad. Nature, 512(7514), 253-253.

Keomany S, Mayxay M, Souvannasing P, Vilayhong C, Stuart BL, Srour L and Newton PN (2007) Toad poisoning in Laos. Am. J. Trop. Med. Hyg. 77(5):.850–853

McClelland, P., Reardon, J. T., Kraus, F., Raxworthy, C. J., & Randrianantoandro, C. (2015). Asian toad eradication feasibility report for Madagascar. Te Anau, New Zealand.

Mecke, S. (2014). Invasive species: review risks before eradicating toads. Nature, 511(7511), 534-534.

Moore M; Fidy JFSN; Edmonds D, 2015. The new toad in town: distribution of the Asian toad, Duttaphrynus melanostictus, in the Toamasina area of eastern Madagascar. Tropical Conservation Science, 8(2):440-455.

http://tropicalconservationscience.mongabay.com/content/v8/tcs\_v8i2\_439-454\_Moore.pdf

Trainor, C.R. 2009. Survey of a population of black-spined toad Bufo melanostictus in Timor-Leste: confirming identity, distribution, abundance and impacts of an invasive and toxic toad. Report by Charles Darwin University to AusAID, contract agreement 52294: 46.

#### **General information**

Berry, P. Y., & Bullock, J. A. (1962). The food of the common Malayan toad, Bufo melanostictus Schneider. Copeia, 736-741. Chicago Church G (1960) The invasion of Bali by Bufo melanostictus. Herpetologica 16(1): 15-21.

Church, G. (1960). The invasion of Bali by Bufo melanostictus. Herpetologica, 16(1), 15-21.

Csurhes, S. 2016. Asian spined toad, Bufo melanosticus. Invasive Animal Risk Assessment. Australia: Department of Agriculture and Fisheries, Biosecurity Queensland.

Dijk PP van, Iskandar D, Lau MWN, Huiqing G, Baorong G, Kuangyang L, Wenhao C, Zhigang Y, Chan B, Dutta S, Inger RF, Manamendra-Arachchi K, Khan MS, 2016. Duttaphrynus melanostictus. IUCN Red List of Threatened Species 2016: e.T54707A86445591. www.iucnredlist.org/

https://www.itis.gov/servlet/SingleRpt/SingleRpt?search\_topic=TSN&search\_value=773557#null http://www.catalogueoflife.org/col/details/species/id/7f7417c5bfd7973da2fd3e19d45bc55b

http://www.iucnredlist.org/details/54707/0

Kaiser H; Lopes Carvalho V; Ceballos J; Freed P; Heacox S; Lester B; Richards SJ; Trainor CR; Sanchez C; O'Shea M, 2001. The herpetofauna of Timor-Leste: a first report. Zookeys, No. 109:19-86. https://doi.org/10.3897/zookeys.109.1439

DAISIE (Delivering Alien Invasive Species Inventories for Europe), 2018.

http://www.europe-aliens.org/speciesFactsheet.do?speciesId=50028#

O'Shea M; Sanchez C; Kathriner A; Mecke S; Lopes Carvalho V; Varela Ribeiro A; Afranio Soares Z; Lemos Araujo L de; Kaiser H, 2015. Herpetological diversity of Timor-Leste: updates and a review of species distributions. Asian Herpetological Research, 6(2):73-131. Seebens H, Blackburn T M, Dyer E E, Genovesi P, Hulme P E, Jeschke J M, Pagad S, Pyšek P, Winter M, Arianoutsou M, Bacher S, Blasius B, Brundu G, Capinha C, Celesti-Grapow L, Dawson W, Dullinger S, Fuentes N, Jäger H, Kartesz J, Kenis M, Kreft H, Kühn I, Lenzner B, Liebhold A, Mosena A (et al), 2017. No saturation in the accumulation of alien species worldwide. Nature Communications. 8 (2), 14435. http://www.nature.com/articles/ncomms14435

Southeast Asian Toad (Duttaphrynus melanostictus) Ecological Risk Screening Summary U.S. Fish & Wildlife Service, January 2021 Wogan GOU; Stuart BL; Iskandar DT; McGuire JA, 2016. Deep genetic structure and ecological divergence in a widespread human commensal toad. Biology Letters, 12(1):20150807. http://rsbl.royalsocietypublishing.org/content/12/1/20150807

Zengeya TA, Wilson JR, 2021. Appendix 2 to "The Status of Biological Invasions and their Management in South Africa in 2019"—the species list., Zenodo. DOI:https://doi.org/10.5281/zenodo.3947659