Ploceus melanocephalus **System:** Terrestrial

Kingdom	Phylum	Class	Order	Family
Animalia	Chordata	Aves	Passeriformes	Ploceidae

Common name Black-headed weaver (English)

Synonym

Similar species

Summary

view this species on IUCN Red List

Principal source:

Compiler:

Review:

Pubblication date:

ALIEN RANGE

[1] BELGIUM
[1] IBERIAN PENINSULA
[1] PORTUGAL
[1] UNITED KINGDOM
[1] GERMANY
[1] ITALY
[1] SPAIN

Red List assessed species 12: LC = 12;

Acrocephalus arundinaceus **LC**
Acrocephalus scirpaceus **LC**
Carduelis carduelis **LC**
Cettia cetti **LC**
Chloris chloris **LC**
Cisticola juncidis **LC**
Locustella luscinioides **LC**
Luscinia megarhynchos **LC**
Passer domesticus **LC**
Passer hispaniolensis **LC**
Remiz pendulinus **LC**
Serinus serinus **LC**

BIBLIOGRAPHY

5 references found for **Ploceus melanocephalus**

Managment information

Grundy, J.P.B., Franco, A.M.A. & Sullivan, M.J.P., (2014). Testing multiple pathways for impacts of the non-native Black-headed Weaver Ploceus melanocephalus on native birds in Iberia in the early phase of invasion. Ibis, 156, pp.355–365.

Sullivan, M.J.P (2014). Predicting the distribution and impacts of non-native birds in the Iberian Peninsula. School of Environmental Sciences, University of East Anglia

Sullivan, M.J.P., Grundy, J. and Franco, A.M.A. (2014), Assessing the impacts of the non-native Black-headed Weaver on native Acrocephalus warblers. Ibis, 156: 231-232. https://doi.org/10.1111/ibi.12104

General information

Monticelli, D. (2008). Finding introduced birds near Lisbon, Portugal. Birding World 21

Sanz-Aguilar A, Carrete M, Edelaar P, Potti J, Tella JL. (2015). The empty temporal niche: breeding phenology differs between coexisting native and invasive birds. Biol Invas. 2015;17:3275–88.