

Homalodisca vitripennis   正體中文

System: Terrestrial

Kingdom	Phylum	Class	Order	Family
Animalia	Arthropoda	Insecta	Hemiptera	Cicadellidae

Common name glassy-winged sharp shooter (English)

Synonym *Homalodisca coagulata*

Similar species *Pheralacerta coagulata*, *Homalodisca liturata*

Summary The glassy-winged sharpshooter (GWSS), *Homalodisca vitripennis* (Germar), is a xylem-feeding leafhopper native to the southeastern United States and regions of northern Mexico. This insect is an important vector of the xylem-limited bacterium, *Xylella fastidiosa*. *X. fastidiosa* is responsible for many economically important diseases including phony peach disease, numerous leaf scold and scorch diseases, variegated citrus chlorosis and Pierce's disease of grape. This insect has been accidentally introduced to California, Arizona and a number of South Pacific islands where it threatens the grape and citrus industries. They lay their eggs inconspicuously below the epidermis of plant leaves which has allowed them to spread to new locations through the nursery trade at an alarming rate.



[view this species on IUCN Red List](#)

Species Description

Glassy-winged sharp shooter (*Homalodisca vitripennis*) adults are about 13-14mm long. They have dark brown bodies with small yellow dots on the head and thorax. The underside of the abdomen has ivory and black markings. They have large translucent smoky-brown wings with red markings. The face and legs are yellow-orange in colour, and the eyes are yellow with dark speckles. Nymphs have a similar body shape to adults but are wingless, gray and have prominent bulging eyes (PaDIL 2006).

Please see PaDIL (Pests and Diseases Image Library) Species Content Page [Bugs: Glassy-Winged Sharp Shooter](#) for high quality diagnostic and overview images.

Notes

The glassy-winged sharp shooter (*Homalodisca vitripennis*) is also referred to as *Homalodisca coagulata* in literature. *Homalodisca vitripennis* is considered a senior synonym of *Tettigonia coagulata* syn. nov. and therefore should be used as the scientific name for the glassy-winged sharpshooter, a major vector of the bacterial Pierce's disease of grapes, phony peach disease, plum leaf scald, and oleander leaf scorch in southern United States and northern Mexico. Please see [Takiya et al. 2006](#) for details.

Lifecycle Stages

There are two generations of *Homalodisca vitripennis* in California per year. Females overwinter as adults and lay eggs in late-winter or early spring. The young wingless nymphs emerge from eggs after about two weeks. Nymphs undergo five molting stages before reaching maturity, during which time they feed on the stems of the host plants. They mature in late spring through summer and lay more eggs. This second generation begins to mature in summer and provides the overwintering adults for the following year (MAF n.d.).

Habitat Description

Glassy-winged sharp shooters (*Homalodisca vitripennis*) are native to the south-eastern part of the USA, where they are found to occur in forest margins. In their introduced range, in California, riparian woodlands in coastal and foothill areas seem to be suitable habitat.

Reproduction

Females lay eggs on host plants below the leaf epidermis once during their two-month (can live much longer when overwintering) adult stage. Oviposition predominantly occurs at night (Tipping *et al.* 2005 in Mizell *et al.* 2008) with 10 to 12 eggs produced at a time. Egg masses vary from 3-28 dependent on past feeding history (Varela *et al.* 2007).

Nutrition

Glassy-winged sharpshooters (*Homalodisca vitripennis*) have a vast host range of over 100 species (ornamental landscape plants, agricultural crops, and natural vegetation; they especially prefer sumac and crepe myrtle). They feed on the xylem (plant tissue that transports water and minerals from roots) of their hosts. Because xylem fluid is nutrient poor *H. vitripennis* displays very high consumption rates (Mizell *et al.* 2008).

Young nymphs feed on the stems of the plant on which they hatched and are known to consume up to 10 times their weight in liquids in one hour. Active stages of the insect must feed constantly to avoid starving to death as adults cannot survive for more than about 4h without food (MAF n.d.). Because xylem is so nutrient poor *H. vitripennis* display host switching which allows them to take advantage of differing nutrient levels between plants. They must respond quickly to changes in xylem fluid content by dispersing to find plants with the highest nutrient levels. Adult glassy-winged sharpshooters are strong fliers and can disperse great distances to search for optimal host plants (Blackmer *et al.* 2003 in Mizell *et al.* 2008).

General Impacts

The glassy-winged sharpshooter is an extremely polyphagous insect, feeding on the xylem fluid of over 100 known species of plant, in at least 37 different families (Alderz, 1980 in Boucias *et al.* 2007). The actual feeding on plants causes little damage; it is the ability of the GWSS to vector the bacterium *Xylella fastidiosa* that causes devastating diseases of many plants which is a major problem (Varela *et al.* 2007).

There are almost 150 known strains of *X. fastidiosa*, which may be benign or cause mild to severe disease symptoms. In grapes one of these strains causes a lethal disease known as Pierce's disease, which is a particular problem in California vineyards as grapes are one of the most economically important crops in California (US \$4.1 billion/year) (Krugner *et al.* 2008). Pierce's disease has been particularly damaging to wine grapes in the Temecula Valley viticulture area, where losses have been as high as 20-30% for some vineyards (Hix 2001 in Byrne and Toscano 2007). Although the disease has been present for many years, it was easily managed because native sharpshooters were poor vectors. The arrival of GWSS, which are far more efficient vectors, has allowed the bacterium to spread much further causing damage to hundreds of hectares of vines (MAF n.d.).

In California *X. fastidiosa* also causes oleander leaf scorch, almond leaf scorch, mulberry leaf scorch, cherry plum leaf scorch and sweet gum dieback. Outside of California other strains of the bacterium cause phony peach disease, plum leaf scald, leaf scorch in sycamore, elm, maple, and oak, and variegated citrus chlorosis (Varela *et al.* 2007).

The glassy-winged sharpshooter is also a problem due to the production of large amounts of white excrement ("sharpshooter rain") which damages cars and other surfaces (Varela *et al.* 2007).

Management Info

The current strategy for containing the problem of GWSS and the disease causing *Xylella fastidiosa* bacterium it transmits is to keep the insect out of new areas. In the United States tighter inspections and regulations of nursery trade in infected areas have been imposed in an attempt to slow down the rapid spread of the pest. The main chemical used to protect *Xylella*-susceptible plants in both commercial agriculture and urban landscapes is imidacloprid, which is registered for home and professional landscape use on nonfood crops. Parasitoid wasps are the most commonly used biocontrol agent against GWSS. Searches for parasitoid wasps have resulted in the collection of several mymarid and trichogrammatid species. In California species released include *Gonatocerus ashmeadi*, *G. fasciatus*, *G. morrilli*, and *G. triguttatus*, and most recently a strain of *Anagrus epos* from Minnesota (Morse *et al.* 2006). Research by Krugner *et al.* (2008) aims to determine host range and efficacy of *A. epos* against GWSS populations.

Please follow this link for more details on the [control options used for the management of the glassy-winged sharpshooter](#).

Pathway

Glassy-winged sharp shooters (*Homalodisca vitripennis*) were probably introduced as eggs on plants.

Principal source:

Compiler: National Biological Information Infrastructure (NBII) & IUCN/SSC Invasive Species Specialist Group (ISSG)

Updates completed with support from Ministry of Agriculture and Forestry (MAF)- Biosecurity New Zealand

Review:

Publication date: 2010-05-28

ALIEN RANGE

[1] CHILE
[7] FRENCH POLYNESIA
[16] UNITED STATES

[1] COOK ISLANDS
[1] HONDURAS

BIBLIOGRAPHY

32 references found for *Homalodisca vitripennis*

Managment information

[Biosecurity New Zealand 2008. Bio-control for Glassy-winged Sharpshooter in Cook Islands.](#)

Summary: Available from: <http://www.biosecurity.govt.nz/biosec/pubs-news/pubs/biosecurity/issue-85/sharpshooter> [Accessed October 10 2009]

[Biosecurity New Zealand. n.d. Glassy-winged sharpshooter factsheet.](#)

Summary: Available from: <http://www.biosecurity.govt.nz/files/pests/glassy-winged-sharpshooter/glassy-winged-sharpshooter-booklet.pdf> [Accessed October 10 2009]

Boucias, D.G., Scharf, D.W., Breaux, S.E., Purcell, D.H. & Mizell, R.E. 2007. Studies on the fungi associated with the glassy-winged sharpshooter *Homalodisca coagulate* with emphasis on a new species *Hirsutella homalodiscae* nom. prov. *BioControl* 52: 231-258.

Castle, S.J. & Naranjo, S.E. 2008. Comparison of sampling methods for determining relative densities of *Homalodisca vitripennis* (Hemiptera: Cicadellidae) on citrus. *Journal of Economic Entomology* 101(1): 226-235.

[Conklin, T. & Mizell, R.F. 2004. Glassy-winged sharpshooter. University of Florida.](#)

Summary: Available from: http://entomology.ifas.ufl.edu/creatures/fruit/glassywinged_sharpshooter.htm [Accessed October 10 2009]

[Grandgirard, Julie.; Mark S. Hoddle, George K. Roderick, Jerome N. Petit, Diana Percy, Rudolph Putoa, Charles Garnier, and Neil Davies.. 2006. Invasion of French Polynesia by the Glassy-Winged Sharpshooter, *Homalodisca coagulata* \(Hemiptera: Cicadellidae\): A New Threat to the South Pacific. *Pacific Science* \(2006\), vol. 60, no. 4:429-438](#)

Summary: Available from: http://muse.jhu.edu/journals/pacific_science/v060/60.4grandgirard.pdf [Accessed 20 March 2008]

Grandgirard, Julie., Mark S. Hoddle, Jerome N. Petit, George K. Roderick and Neil Davies., 2008. Engineering an invasion: classical biological control of the glassy-winged sharpshooter, *Homalodisca vitripennis*, by the egg parasitoid *Gonatocerus ashmeadi* in Tahitiand Moorea, French Polynesia. *Biological Invasions* (2008) 10:135-148

Summary: En 2004, un programme de contrôle biologique classique contre la Cicadelle pisseuse a été lancé en Polynésie française. Cet article présente les premiers résultats de ce programme un an après le lâcher du parasitoïde en 2005.

Global Invasive Species Database (GISD) 2025. Species profile *Homalodisca vitripennis*. Available from: <https://www.iucngisd.org/gisd/species.php?sc=240> [Accessed 09 May 2025]

Hall, D.G. & Hunter, W.B. 2008. Populations of sharpshooters in two citrus groves in eastcentral Florida as indicated by yellow sticky card traps. *Florida Entomologist* 91(3): 488-490.

Hoddle M.S., Grandgirard J., Petit J., Roderick G.K., Davies N., 2006. Glassy-winged sharpshooter Ko ed - First round - in French Polynesia. *Biocontrol News and Information* 27(3), 47N-62N

Summary: Cet article présente les premiers résultats du programme de lutte biologique contre la Cicadelle pisseuse en Polynésie française. Les résultats montrent que le parasitoïde a rapidement colonisé toute l'île de Tahiti et que l'abondance de la Cicadelle pisseuse a chuté de plus de 90%.

Hunnicut, L. E.; J. Mozoruk, W. B. Hunter, J. M. Crosslin, R. D. Cave, C. A. Powell., 2008. Prevalence and natural host range of *Homalodisca coagulata* virus-1 (HoCV-1). *Arch Virol* (2008) 153: 61-67

Hunnicut, L.E., Mozoruk, J., Hunter, W.B., Crosslin, J.M., Cave, R.D. & Powell, C.A. 2008. Prevalence and natural host range of *Homalodisca coagulata* virus-1 (HoCV-1). *Archives of Virology* 153: 61-67.

[Introduced Species Summary Project., 2002. Glassy-winged Sharpshooter, Xylophagous Leafhopper. Andria Cimino and Edited by James A. Danoff-Burg, Columbia University \(February 2002\).](#)

Summary: Report on taxonomy, description, distribution, mode of introduction, effects of introduction, and control methods.

Available from: http://www.columbia.edu/itc/cerc/danoff-burg/invasion_bio/inv_spp_summ/Homalodisca_coagulata.html#Distribution [Accessed December 2002]

Kanga, L.H.B., Jones, W.A., Humber, R.A. & Boyd, D.W. 2004. Fungal pathogens of the glassy-winged sharpshooter *Homalodisca coagulata* (Homoptera: Cicadellidae). *Florida Entomologist* 87(2): 225-228.

Krugner, R., Johnson, M.W., Groves, R.L. & Morse, J.G. 2008. Host specificity of *Anagrus epos*: a potential biological control agent of *Homalodisca vitripennis*. *BioControl* 53: 439-449.

Lauziere, I. & Elzen, G. 2007. Effect of formulated insecticides on *Homalodisca vitripennis* (Germar) (Hemiptera: Cicadellidae) and its parasitoid *Gonatocerus ashmeadi girault* (Hymenoptera: Mymaridae). *Journal of Entomological Science* 42(1): 11-19.

Lutte biologique contre la Cicadelle pisseuse en Polynésie française. Site internet

Summary: Site internet sur la lutte biologique contre la Cicadelle pisseuse en Polynésie française. Ce site contient de nombreuses informations et des articles en téléchargement.

Miller, T.A. 2007. Applying insect transgenic technology: scientific and regulatory experiences. *Entomological Research* 37: 67-75.

[Ministry of Agriculture and Forestry \(MAF\), Media Release 4 April 2007. Cook Island imports under extra scrutiny after insect pest detected](#)

Summary: Available from: <http://www.maf.govt.nz/mafnet/press/070404gwss.htm> [Accessed 20 March 2008]

Morse, J.G., Morgan, D.J.W., Lytle, J.M. 2006. Seasonal population dynamics of glassy-winged sharpshooter egg parasitoids: variability across sites and host plants. In: Esser T, Tariq MA, Medeiros R, Mochel M, Veling S (eds) *Proceedings, 2006 Pierce's disease research symposium*, San Diego, CA, 27-29 November 2006. Copeland Printing, Sacramento, CA, pp 92-94.

Pacific Pest Info Newsletter. Published by the Secretariat of the Pacific Community: Plant Protection Service, Private Mail Bag, Suva, Fiji Islands. Tel: (679) 3370-733; Fax: (679) 3370-021.

Petit, Jerome N.; Mark S. Hoddle, Julie Grandgirard, George K. Roderick and Neil Davies., 2007. Invasion dynamics of the glassy-winged sharpshooter *Homalodisca vitripennis* (Germar) (Hemiptera: Cicadellidae) in French Polynesia. *Biological Invasions*. DOI 10.1007/s10530-007-9172-3

Summary: Les conclusions de cet article suggèrent que 1) l'abondance de la Cicadelle pisseuse est fortement corrélée au niveau d'urbanisation, (2) la cicadelle pisseuse peut montrer un effet Allee pendant la phase la plus précoce de l'invasion, et (3) la dynamique d'invasion est marquée par les moyens de déplacements humains, rapides et sur de longues distances.

Ramirez, J.L., Perring, T.M. & Miller, T.A. 2008. Fate of a genetically modified bacterium in foregut of glassy-winged sharpshooter (Hemiptera: Cicadellidae). *Journal of Economic Entomology* 101(5): 1519-1525.

Sisterson, M.S., Yacoub, R., Montez, G., Grafton-Cardwell, E.E. & Groves, R.L. 2008. Distribution and management of citrus in California: implications for management of glassy-winged sharpshooter.

University of California, Riverside. The Glassy-Winged Sharpshooter.

Summary: Report on distribution in California, description, biology, details about Pierce's disease and its spread, and potential management plans.

[University of California., undated. Integrated Pest Management of the Glassy-winged Sharpshooter and the Diseases it Transmits: Frequently Asked Questions. University of California Glassy-winged Sharpshooter Workgroup.](#)

Summary: Reports on description, effects of introduction, distribution, habitat, biological management, description of and spread of Pierce's Disease, and management of spread.

Available from: <http://gwss.ucanr.org/faq.html#6> [Accessed December 2002]

[USDA. 2005. News and Events: Scientists Seeking Biocontrols Against Sharpshooters. Alfredo Flores, Public Affairs Specialist, Agricultural Research Service of the U.S. Department of Agriculture.](#)

Summary: Available from: <http://www.ars.usda.gov/is/pr/2005/051020.htm> [Accessed January 2006]

[Varela, L.G., Hashim-Buckey, J.M. & Phillips, P.A. 2007. Glassy-winged sharpshooter. Pest Notes Publication 7492. University of California Agriculture and Natural Resources.](#)

Summary: Available from: <http://www.ipm.ucdavis.edu/PMG/PESTNOTES/pn7492.html> [Accessed October 10 2009]

[Walker, K. 2006. Glassy-Winged Sharp Shooter \(*Homalodisca vitripennis*\) Pest and Diseases Image Library. Updated on 10/08/2006 10:05:58 AM.](#)

Summary: PaDIL (Pests and Diseases Image Library) is a Commonwealth Government initiative, developed and built by Museum Victoria's Online Publishing Team, with support provided by DAFF (Department of Agriculture, Fisheries and Forestry) and PHA (Plant Health Australia), a non-profit public company. Project partners also include Museum Victoria, the Western Australian Department of Agriculture and the Queensland University of Technology. The aim of the project is: 1) Production of high quality images showing primarily exotic targeted organisms of plant health concern to Australia. 2) Assist with plant health diagnostics in all areas, from initial to high level. 3) Capacity building for diagnostics in plant health, including linkage developments between training and research organisations. 4) Create and use educational tools for training undergraduates/postgraduates. 5) Engender public awareness about plant health concerns in Australia. PaDIL is available from : <http://www.padil.gov.au/aboutOverview.aspx>, this page is available from: <http://www.padil.gov.au/viewPestDiagnosticImages.aspx?id=173> [Accessed 6 October 2006]

General information

[European and Mediterranean Plant Protection Organization \(EPPO\), 2002. *Homalodisca coagulata*. European and Mediterranean Plant Protection Organization. \(July, 2002\).](#)

Summary: Report on description, biology, and effects on California's grapevines and other crops.

Available from: http://www.eppo.org/QUARANTINE/insects/Homalodisca_coagulata/HOMLTR-a.htm [Accessed 27 May 2010]

Grandgirard, J., Hoddle, M.S., Petit, J., Roderick, G.K. & Davies, N. 2007. Classical biological control of the glassy-winged sharpshooter, *Homalodisca vitripennis*, by the egg parasitoid *Gonatocerus ashmeadi* in the Society Marquesas and the Austral archipelagos of French Polynesia. *Biological Control* 48: 155-163.

Mizell, R.F., Tipping, C., Andersen, P.C., Brodbeck, B.V., Hunter, W.B. & Northfield, T. 2008. Behavioral model for *Homalodisca vitripennis* (Hemiptera: Cicadellidae): optimization of host plant utilization and management implications. *Environmental Entomology* 37(5): 1049-1062.

[Takiya, D.M., Stuart H.M., and Rodney R.C., 2006. Validity of *Homalodisca* and of *H. vitripennis* as the Name for Glassy- Winged Sharpshooter \(Hemiptera: Cicadellidae: Cicadellinae\) Ann. Entomol. Soc. Am. 99\(4\): 648-655 \(2006\)](#)