

FULL ACCOUNT FOR: Wisteria floribunda

Wisteria floribunda 简体中文 正體中文

| Kingdom | Phylum        | Class         | Order   | Family   |
|---------|---------------|---------------|---------|----------|
| Plantae | Magnoliophyta | Magnoliopsida | Fabales | Fabaceae |

Common name

Japanese Wisteria (English, United States)

**Synonym** 

Dolichos japonicus, Spreng. 1826

Kraunhia brachybotrys, (Siebold & Zucc.) Greene 1892

Glycine floribunda, Willd. 1802

Kraunhia floribunda, (Willd.) Taub. forma albiflora Makino 1911

Kraunhia floribunda, (Willd.) Taub. var. brachybotrys (Siebold & Zucc.) Makino

1911

Kraunhia floribunda, (Willd.) Taub. 1894

Kraunhia floribunda , (Willd.) Taub. var. typica Makino 1911 Kraunhia floribunda, (Willd.) Taub. var. pleniflora Makino 1911 Kraunhia sinensis, (Sims) Makino forma albiflora Makino 1910 Kraunhia sinensis, (Sims) Makino var. pleniflora Makino 1910

Kraunhia sinensis, (Sims) Makino var. brachybotrys (Siebold & Zucc.) Makino

Kraunhia sinensis, (Sims) Makino var. floribunda (Willd.) Makino 1910

Millettia floribunda, (Willd.) Matsum. 1902

Phaseoloides brachybotrys, (Siebold & Zucc.) Kuntze 1891

Phaseoloides floribunda, (Willd.) Kuntze 1891 Rehsonia floribunda, (Willd.) Stritch 1984 Wisteria brachybotrys, Siebold & Zucc. 1839

Wisteria chinensis, DC. var. multijuga (Van Houtte) Hook.f. 1897

Wisteria chinensis, DC. var. macrobotrys (Siebold ex Neubert) Lavallee 1877

Wisteria chinensis, DC. var. flore-plena (Carrirre) W.Mill. 1902

Wisteria floribunda , (Willd.) DC. forma rosea (Bean) Rehder & E.H.Wilson 1916 Wisteria floribunda, (Willd.) DC. forma macrobotrys (Siebold ex Neubert) Rehder & E.H.Wilson 1916

Wisteria floribunda, (Willd.) DC. forma variegata (G.Nicholson) Rehder & E.H.Wilson 1916

Wisteria floribunda, (Willd.) DC. forma alba (Carrirre) Rehder & E.H.Wilson 1916

Wisteria macrobotrys, Siebold ex Neubert 1870

Wisteria multijuga , Van Houtte var. rosea Bean 1914

Wisteria multijuga, Van Houtte 1874

Wisteria polystachya , K.Koch forma alba (Carri@re) Zabel 1903

Wisteria multijuga , Van Houtte var. alba Carri@re 1891

Wisteria multijuga , Van Houtte var. variegata G.Nicholson 1887

Wisteria polystachya , K.Koch forma variegata (G.Nicholson) Zabel 1903 Wisteria polystachya , K.Koch forma multijuga (Van Houtte) Zabel 1903 Wisteria sinensis, (Sims) Sweet var. violaceo-plena C.K.Schneid. 1907 Wisteria sinensis, (Sims) Sweet forma violaceo-plena Rehder & E.H.Wilson

1916

Similar species

Wisteria sinensis. Wisteria frutescens

Summary

In its alien range, Wisteria floribunda is still used as an ornamental and often escapes from landscapes and becomes invasive in natural ecosystems. Wisteria floribunda infests forest edges and disturbed areas, including riparian zones and tolerates shade and a variety of soil types. In riparian areas, Wisteria floribunda spreads downstream as seeds float to new locations.

**System:** Terrestrial



FULL ACCOUNT FOR: Wisteria floribunda



view this species on IUCN Red List

# **Species Description**

According to Martin (2002), *Wisteria floribunda* is a perennial vine that can live for 50 years or more and can grow up to 38cm in diameter. Leaves are alternate and pinnately compound; they are up to 30cm long and consist of 13-19 leaflets. Flowers hang in clusters that sometimes exceed 40cm in length. Flower colour is usually blue-violet but other cultivars (which may be escaped plants) are white, purple, pink, and lavender. Flowers are usually produced from April to May in the United States. Seedpods are 10-15cm in length, hairy, brown, narrow at the base, and constricted between seeds. *W. floribunda* twines clockwise around host plants.

### Notes

The seeds and pods of *Wisteria floribunda* are toxic if ingested and may cause nausea, vomiting, stomach pains, and diarrhea (Martin, 2002). *Wisteria floribunda* is not as widespread in the United States as its brother vine, *W. sinensis*, which has caused problems throughout the eastern U.S.

### Uses

Wisteria floribunda has been commonly used as an ornamental since 1830 when it was first introduced (Remaley, 1999). Martin (2002) notes that it has been grown extensively in the United States for ornamental uses on porches, gazebos, walls, and gardens. Many alternatives to this vine exist, including American wisteria (Wisteria frutescens), Trumpet creeper (Campsis radicans), Trumpet honeysuckle (Lonicera sempervirens), Dutchman's pipe/pipevine (Aristolochia macrophylla), and Crossvine (Bignonia capreolata).

# **Habitat Description**

Typically, Wisteria floribunda infests forest edges and disturbed areas (Martin, 2002). Remaley (1999) notes that it grows best in full sun but is shade-tolerant. W. floribunda can tolerate a variety of soil and moisture types but it prefers loamy, deep, well drained soils.

# Reproduction

According to Remaley (1999), vegetative reproduction is the primary means of expansion for *Wisteria floribunda*. In its preferred habitat, however, seeds may be produced, and in riparian areas they may be carried downstream for long distances.

# **General Impacts**

According to Remaley (1999), native shrubs are overtaken by *W. floribunda* through strangling and shading. Even larger trees can be killed by this vine, causing large gaps in the canopy when they fall; this open canopy furthers the growth of *W. floribunda*. This aggressive vine may form dense thickets allowing little else to grow.



FULL ACCOUNT FOR: Wisteria floribunda

# **Management Info**

<u>Physical</u>: According to Martin (2002), current management approaches consist of mechanical and chemical methods. Mechanical methods should be used for small populations or where herbicides could damage desirable species. When cutting vines, cut close to the root collar to discontinue growth of existing vines and reduce seed production. *W. floribunda* will resprout, so it is recommended that the vine be cut repeatedly every two weeks from early in the growing season to autumn. Vines should be removed because they may continue to grow and girdle the host plant. Try to remove the entire plant (including roots) and dispose of all parts because any plant parts left can resprout.

\r\n<u>Chemical</u>: Cut-stump herbicide applications should be used where there are large stands of established vines or where desirable plants occur and could be affected by foliar spray. Cut the vine close to the ground and apply glyphosate or triclopyr (25% solutions in water) to the cut area. If resprouting occurs retreatment may be necessary. This treatment is not effective if the ground is frozen. Foliar sprays should be used where mechanical controls would be disruptive and cut-stump methods are impractical, but additional precautions should be taken not to harm non-target species. Spray the foliage thoroughly, but do not apply so much that it drips off. Application may be more effective in warmer temperatures (above 15-18 C) because translocation is slower in cooler weather. Triclopyr is specific for control of broadleaved plants and may be beneficial if protection of valuable native grasses is of concern. Glyphosate is non-selective and should be used with care. Chlopyralid targets aster, buckwheat, and the pea family. However, chlopyralid can seep into groundwater in sandy and limestone soil types. Picloram may provide control in areas where desirable vegetation is not present.

# **Pathway**

W. floribunda is a popular ornamental (Martin, 2002).

Principal source: Exotic Wisterias (Remaley, 1999)

Compiler: National Biological Information Infrastructure (NBII) & IUCN/SSC Invasive Species Specialist Group

(ISSG)

Review: Kristine Johnson Supervisory Forester Great Smoky Mountains National Park. USA

Pubblication date: 2005-01-24

**ALIEN RANGE** 

[3] UNITED STATES

## **BIBLIOGRAPHY**

4 references found for Wisteria floribunda

### Managment information

Martin, Tunyalee. 2002. Weed Notes: Wisteria sinensis (Chinese Wisteria) Wisteria floribunda (Japanese Wisteria) . The Nature Conservancy: Wildland Invasive Species Team [Online database].

**Summary:** Detailed description, distribution, biology, similar species description, alternative ornamental plants, and control measures used. Available from: http://tncweeds.ucdavis.edu/ [Accessed 16 April 2003].

### **General information**

ITIS (Integrated Taxonomic Information System), 2005. Online Database Wisteria floribunda

**Summary:** An online database that provides taxonomic information, common names, synonyms and geographical jurisdiction of a species. In addition links are provided to retrieve biological records and collection information from the Global Biodiversity Information Facility (GBIF) Data Portal and bioscience articles from BioOne journals.

Available from:

 $\label{lem:http://www.cbif.gc.ca/pls/itisca/taxastep?king=every&p_action=containing\&taxa=Wisteria+floribunda&p_format=&p_ifx=plglt&p_lang=[Accessed March 2005]$ 

Plants & Japan, Masashi Yamaguchi

Summary: Japanese plants.

Available from: http://homepage3.nifty.com/plantsandjapan/index.html [Accessed 24 September 2004]



FULL ACCOUNT FOR: Wisteria floribunda

Wunderlin, Richard and Hansen, Bruce. 2002. Wisteria sinensis Institute of Systematic Botany: Atlas of Florida Vascular Plants [Online database].

**Summary:** Detailed taxonomy and Florida distribution.