

GLOBAL INVASIVE SPECIES DATABASE

Sagittaria sagittifolia 简体中文 正體中文

System: Terrestrial

Kingdom	Phylum	Class	Order	Family
Plantae	Magnoliophyta	Liliopsida	Alismatales	Alismataceae
Common name	suokiu (English), shui p'Ing (English), chieh ku (English), giant arrowhead (English), pijlkruid (English), pai ti li (English), chien tao ts'ao (English), flecha de agua (Spanish), t'zu ku (English), sagit�ria (Portuguese), espadana (Portuguese), wapatoo (English), Pfeilkraut (German), fl�che d'eau (French), yen wei ts'ao (English), old world arrowhead (English), saeta de agua (Spanish), arrowhead (English), Hawaii arrowhead (English)			
Synonym	,ittaria japonica (Hort.) Sagittaria sagittifolia , var. edulis Siebold ex Miq. Sagittaria sagittifolia , var. leucopetala Miq. Sagittaria sinensis , Sims Sagittaria trifolia , L. Sagittaria trifolia , var. edulis (Siebold ex Miq.) Sagittaria trifolia , var. sinensis (Sims) Makino			
Similar species				
Summary	Sagittaria sagittifolia is a very hardy aquatic plant that has become a general nuisance in the crops' irrigation systems, drains and waterways around the globe.			
CEP	view this species on IUCN Red List			

Species Description

Sagittaria sagittifolia is a herbaceous aquatic perennial that reaches sizes of 1m by 0.5m. It is a hardy species that is not frost tender. The leaves are borne on triangular stalks that vary in length with the depth of the water in which the plant is growing. They do not lie on the water but stand boldly above it. They are large and arrow-shaped and very glossy. The early, submerged leaves are ribbon like. The flower-stem rises directly from the root and bears several rings of buds and blossoms, three in each ring or whorl, and each flower composed of three outer sepals and three large, pure white petals, with a purple blotch at their base. The upper flowers are stamen bearing; the lower ones generally contain the seed vessels only. *S. sagittifolia*'s tubers are walnut sized and grow just below the mud surface, produced on creeping runners (Grieve, 2005; National Plant Pest Accord, 2001; and Plants for a Future, 2000).

Lifecycle Stages

North American region: *S. sagittifolia* flowers in mid-summer, and the seeds ripen through the fall. The flowers are monoecious (individual flowers are either male or female, but both sexes can be found on the same plant) and are pollinated by insects (Plants for a Future, 2000).



GLOBAL INVASIVE SPECIES DATABASE

FULL ACCOUNT FOR: Sagittaria sagittifolia

Uses

Plants for a Future (2000) offers a variety of uses both edible and medicinal for *S. sagittifolia* . The leaves and roots of *S. sagittifolia* are edible. The root can be cooked and is apparently excellent when roasted and likened to potatoes. The tubers are starchy with a distinct flavour, but should not be eaten raw. The leaves and young stems can also be cooked but are apparently somewhat acrid. The plant also has medicinal properties as an antiscorbutic; diuretic, and galactofuge but may induce premature birth. The authors post a disclaimer for all edible and medical knowledge: \"We are not experts on the medicinal uses of plants and much of the information has been taken from other sources. You should talk to someone who knows what they are on about before using any of these plants. Plants For A Future can not take any responsibility for any adverse effects from the use of plants\" (Plants for a Future, 2000).

Habitat Description

Sagittaria sagittifolia can inhabit ponds, canals and slow flowing water on muddy sub-strata in water up to 45cm deep, in acid or calcareous conditions (Plants for a Future, 2000). Scher (UNDATED) adds that *S. sagittifolia* can be found in Sub-arctic to tropical environments, in quiet, shallow, standing waters, including swamps, reservoirs, rice paddies, river banks, bays.

Reproduction

Reproduction is by achenes and vegetatively by whole, immature plants and underground tubers. Seeds float easily and can be carried long distances (Scher, UNDATED).

General Impacts

The Nature Conservancy (2005) explains that *S. sagittifolia* is a general nuisance in the crops' irrigation systems, drains and waterways of more than 50 countries. This invasive potential stems from its remarkable ability to adapt, both in form and physiology to a variety of habitats.

Principal source: Plants for a Future, 2000. Sagittaria sagittifolia

Compiler: National Biological Information Infrastructure (NBII) & IUCN/SSC Invasive Species Specialist Group (ISSG) with support from the Terrestrial and Freshwater Biodiversity Information System (TFBIS) Programme (Copyright statement)

Review: Dr. Surrey Jacobs Principal Research Scientist Royal Botanic Gardens Sydney NSW, Australia

Pubblication date: 2006-11-29

ALIEN RANGE

[1] ARGENTINA[1] CUBA[5] NEW ZEALAND

[1] AUSTRALIA[1] MEXICO[2] UNITED STATES

BIBLIOGRAPHY

18 references found for Sagittaria sagittifolia

Managment information

Champion, P. Clayton, J. and Rowe, D. 2002. Alien Invaders Lake Managers Handbook. Ministry for the Environment. **Summary:** Available from: http://www.mfe.govt.nz/publications/water/Im-alien-invaders-jun02.pdf [Accessed 3 February 2005] Champion, P.D.; Clayton, J.S. 2000. Border control for potential aquatic weeds. Stage 1. Weed risk model. Science for Conservation 141. . **Summary:** This report is the first stage in a three-stage development of a Border Control Programme for aquatic plants that have the potential to become ecological weeds in New Zealand.

Available from: http://www.doc.govt.nz/upload/documents/science-and-technical/sfc141.pdf [Accessed 13 June 2007]



GLOBAL INVASIVE SPECIES DATABASE

FULL ACCOUNT FOR: Sagittaria sagittifolia

Champion, P.D.; Clayton, J.S. 2001. Border control for potential aquatic weeds. Stage 2. Weed risk assessment. Science for Conservation 185. 30 p.

Summary: This report is the second stage in the development of a Border Control Programme for aquatic plants that have the potential to become ecological weeds in New Zealand. Importers and traders in aquatic plants were surveyed to identify the plant species known or likely to be present in New Zealand. The Aquatic Plant Weed Risk Assessment Model was used to help assess the level of risk posed by these species. The report presents evidence of the various entry pathways and considers the impact that new invasive aquatic weed species may have on vulnerable native aquatic species and communities.

Available from: http://www.doc.govt.nz/upload/documents/science-and-technical/SFC185.pdf [Accessed 13 June 2007] National Pest Plant Accord, 2001. Biosecurity New Zealand.

Summary: The National Pest Plant Accord is a cooperative agreement between regional councils and government departments with biosecurity responsibilities. Under the accord, regional councils will undertake surveillance to prevent the commercial sale and/or distribution of an agreed list of pest plants.

Available from: http://www.biosecurity.govt.nz/pests-diseases/plants/accord.htm [Accessed 11 August 2005] Royal New Zealand Institute of Horticulture (RNZIH), 2005. Arrowhead Sagittaria sagittifolia Summary: Available from: http://www.rnzih.org.nz/pages/nppa_011.pdf [Accessed 1 October 2005]

General information

Champion, Paul., Undated. Managing tomorrow s weeds today - a risk assessment approach to aquatic weed menagement. NIWA (National Institute of Water & Atmospheric Research).

Summary: Available from: http://www.niwa.co.nz/pubs/wa/12-3/weed [Accessed 11 August 2005]

Combroux, C. S. I., G. Bornette, and C. Amorosa. 2002. Plant Regenerative Strategies after a Major Disturbance: The Case of a Riverine Wetland Restoration. Wetlands: Vol. 22, No. 2, pp. 234-246.

Dhora, D., S. Bequiraj, and D. Dhora. 2001. Report on biodiversity of River Buna. Association for Protection of Aquatic Wildlife of Albania Ecological Association Kalimera, Ulcinj, Montenegro.

Dorken, M. E., and S. C. H. Barrett. 2003. Gender plasticity in *Sagittaria sagittifolia* (Alismataceae), a monoecious aquatic species. Plant Systematics and Evolution 237(1-2):99 - 106.

Global Biodiversity Information Facility (GBIF), 2010. Sagittaria sagittifolia L..

Summary: Available from: http://www.gbif.net/species/13751042/ [Accessed 15 June 2010]

Grieve, M. 2005. Botanical.com: A Modern Herbal.

Summary: Available from: http://www.botanical.com/botanical/mgmh/a/arrow063.html [Accessed 11 August 2005] Hulina, N. 1998. Rare, endangered or vulnerable plants and neophytes in a drainage system in Croatia. Natura Croatica [Nat. Croat.]. Vol. 7, no. 4, pp. 279-289. 31 Dec 1998.

ITIS (Integrated Taxonomic Information System), 2005. Online Database Sagittaria sagittifolia

Summary: An online database that provides taxonomic information, common names, synonyms and geographical jurisdiction of a species. In addition links are provided to retrieve biological records and collection information from the Global Biodiversity Information Facility (GBIF) Data Portal and bioscience articles from BioOne journals. Available from:

http://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=38931 [Accessed 11 August 2005] Pallewatta, N., J.K. Reaser, and A.T. Gutierrez. (eds.). 2003. Invasive Alien Species in South-Southeast Asia: National Reports & Directory of Resources. Global Invasive Species Programme, Cape Town, South Africa.

Plants for a Future. 2000. Sagittaria sagittifolia. Plants For A Future - Species Database.

Summary: Available from: http://www.ibiblio.org/pfaf/cgi-bin/arr_html?Sagittaria+sagittifolia&CAN=COMIND [Accessed 11 August 2005] Scher, J. Undated. Federal Noxious Weed Disseminules of the U. S.

Summary: Available from: http://www.lucidcentral.org/keys/FNW/FNW%20seeds/html/fact%20sheets/Sagittaria%20sagittifolia.htm [Accessed 11 August 2005]

Slovak National Čouncil. 1997. National Strategy of Biodiversity Protection in the Slovak Republic. Government of the Slovak Republic. The Nature Conservancy. 2005. Rod Randall s Big Weed List.

Summary: Available from: http://tncweeds.ucdavis.edu/global/australia/aca.html [Accessed 11 August 2005]