

GLOBAL INVASIVE SPECIES DATABASE

EICAT profile: Tyto novaehollandiae

MV (Massive) Tyto novaehollandiae

Date assessed 2020-05-19 Year published 2021

Eicat category

MV (Massive)

Justification for EICAT assessment

Several species of owl were introduced to Lord Howe Island in the 1920s in an attempt to control black rats (Rattus rattus). Experts beleive that the Lord Howe southern boobook (Ninox novaeseelandiae albaria) disappeared from the island in the 1950s, and that the introduced Australian masked owl probably preyed on the boobook, which may have caused its extinction (NSW Office of Environment & Heritage, 2020). Owls are known to prey on other owl species (see Mikkola, 1976), and the Australian masked owl is a large powerful owl - the native boobook owl was a relatively small species by comparison. Indeed, the Australian masked owl is known to prey on other boobook species in its native range (NSW Office of Environment & Heritage, 2020). However, there is no direct evidence of Australian masked owls preying on the Lord Howe southern boobook on Lord Howe Island. Other possible causes of the boobook's extinction include competition from other introduced owl species such as the barn owl (Tyto alba), and predation by black rats.

Confidence rating

Low

Mechanism(s) of maximum impact

Predation

Countries of most severe

Australia

impact

Description of impact

The Australian masked owl is believed to have preyed on the Lord Howe southern boobook on Lord Howe Island. This predation, probably in combination with competition from other introduced owl species and predation by black rats, is likely to have caused the extinction of the boobook.

Assessor Thomas Evans

Contributors

Reviewers EICAT authority

Recommended citation Thomas Evans. (2025). Tyto novaehollandiae . IUCN Environmental Impact Classification for Alien Taxa



GLOBAL INVASIVE SPECIES DATABASE

EICAT profile: Tyto novaehollandiae

